

side of his head, which he had not done for long previous. The ulcer soon began to heal and after two months and a half of treatment was almost closed over and the infiltrated tissue entirely gone away. At the end of three months treatment it was scabbed over and the man allowed to cross the Atlantic, returning a month later with the ulcer still closed over and no appearance of a recurrence.

A case of *tubercular lupus* of eight years' standing and now with three patches on the face, which had been frequently operated upon without improvement, was treated in the beginning of June, 1901. Eight treatments of X-rays were administered for five minutes every other day at a distance from the tube of 12 inches, with a current of 2 ampères at 110 volts through a Caldwell break. My absence in England then made a break in the treatment until October, but the patient continued to gain. The treatment was renewed every other day, as before, and gradually increased to five minutes every day, and finally to ten minutes every day. In January and February the treatment had to be stopped in consequence of an attack of measles. After the measles the case was seen to be much improved and after a lapse of nearly two months, treatment as before was again renewed, but the improvement did not seem to be so rapid as formerly, therefore a high frequency current was applied with improvement again. This was continued for a month and then it seemed to lose its power to improve the condition and recourse was had to the arc lamp of Lortet and Gnaud's modification of Finzen's light, with the addition of adrenalin as a means of increasing the anæmic condition of the skin. Success followed this, and by the beginning of July the patient was sent away to the country returning well at the end of a month of country life.

A case of *Rodent Ulcer* of the nose near the tip and extending for one inch up towards the bridge of the nose, with a diameter from right to left of one and a quarter inches, a regular cauliflower appearance, and with everted blue edges, had been under treatment in the ordinary manner for six years with no success. The surface of the ulcer was exposed to X-rays at a distance of six inches, the rest of the face and the body being protected by lead sheet, for ten minutes every day, for eleven days, at the end of which time the healing process had been well established. The ulcer healed under antiseptic dressings in 21 days, but a small opening through into the nostril on one side is still left, and this, when the new tissue has had time to become sound, will probably be repaired by paring the edges and bringing them together.

Another case of *Rodent Ulcer* with a hardened and thick base of