

duces perfect therapeutic results, such as I have noted with Ergo-Apiol (Smith) in insane women, may be expected to perform the same services even more promptly in the average case of amenorrhea as met with in ordinary family practice. This is proved conclusively in the numerous cases reported by various observers who employed Ergo-Apiol (Smith) in menstrual disorders, and a partial list of whose publications appear in the annexed bibliography. Ergo-Apiol in the shape of capsules administered three times daily in doses of one or two beginning a little before the expected menses, and continuing through the period, has proven the most efficient, prompt, safe, and pleasant emmenagogue that I have ever employed. My experience with the drug was such as to lead me to adopt it as a routine treatment in amenorrhea.

BIBLIOGRAPHY.

1. Levasseur, F., Clinical Observations on Amenorrhea and Dysmenorrhea. (*International Journal of Surgery*, July, 1901.)
2. Ergo-Apiol (Smith), as an Emmenagogue and Oxytocic, by M. A. Auerbach, M.D., New York. (*Obstetrics*, June, 1901.)
3. Mallette, E. Alfred, Philadelphia (Clinical Report on Ergo-Apiol, 1901.)
4. Shepard, Charles H., Physician to Lincoln Hospital, Durham, N.C., Ergo-Apiol (Smith) in Diseases of the Female. (*Medical Examiner*, July, 1902.)
5. Canan, C. W. Ergo-Apiol (Smith), Its Therapeutical Indications; With Clinical Notes. (*Medical Fortnightly*, July, 1901.)
6. Black, James A. Clinical Report on Ergo-Apiol (Smith), Morganza, Pa., May 15, 1902.
7. Baker, Clayton H., M.B., F.R.S. (London). Disorders of the Uterine Function (*Clinical Report*, New York, 1902.)

—From the *Brooklyn Med. Jour.*, June, 1903.