

Adrenalin Ampoules



AMPOULES OF

Adrenalin Chloride Solution, 1:10,000

(1 Cc. GLASS CONTAINERS)

FOR HYPODERMATIC INJECTION.

The most powerful of astringents and hemostatics, Adrenalin Chloride Solution lends itself to many practical uses. Since the time of its introduction it has been marketed in ounce vials, and of the strength of 1:1000. Experience has shown, however, that a weaker solution is much more frequently required than the "full strength"; and while it is generally an easy matter to dilute with water or normal saline solution, in certain emergencies an already fully diluted preparation is to be preferred. For hypodermatic injection the small hermetically-sealed package will be found of great convenience.

SOME INDICATIONS FOR THE HYPODERMATIC USE OF ADRENALIN CHLORIDE SOLUTION.

Shock	Postpartum hemorrhage
Cardiac and vasomotor asthenia	Asthma
Inaccessible hemorrhages	Osteomalacia
	Whooping-cough

DIRECTIONS—Break off the neck of the ampoule at the file-mark, as shown in the illustration. Use an ordinary hypodermatic syringe. Insert the point of the needle behind the shoulder of the ampoule—not to the bottom. (See cut.) Elevate the bottom of the ampoule as the piston of the syringe is withdrawn, and the contents can be removed to the last drop.

Marketed in boxes of 1 dozen.

KEEP A PACKAGE IN YOUR EMERGENCY CASE.

PARKE, DAVIS & COMPANY

LABORATORIES: Detroit, Mich.; Walkerville, Ont.; Hounslow, Eng.
BRANCHES: New York, Chicago, St. Louis, Boston, Baltimore, New Orleans, Kansas City,
Minneapolis; London, Eng.; Montreal, Que.; Sydney, N.S.W.; St. Petersburg, Russia;
Bombay, India; Tokio, Japan; Buenos Aires, Argentina.