

HEADACHE POWDERS.

Directions for use as found on the label.	Remarks.	RESULTS OF ANALYSIS.					No. of Sample.	Remarks and Opinion of the Chief Analyst.
		Weights of Constituents of one powder, grains.	Acetanilide = Antifebrin, grains.	Phenacetin, grains.	Phenazone = Antipyryn, grains.			
	Powder No. 1	4.7	2.2	None.	None.	47588	Curative powers claimed. Meets requirements of P. or P. Act. Acetanilide declared.	
	" No. 2	4.9	2.3	"	"			
	Powder No. 1	12.1	5.0	None.	None.	47589	Curative powers claimed. Acetanilide declared.	
	" No. 2	10.9	4.4	"	"			
Directions:—For Headache, one wafer which may be repeated in thirty minutes if necessary. Dip the wafer in water until thoroughly soft, then place on the tongue and swallow with a little water, at least three hours should elapse before another dose is taken. A sedlitz powder or other mild laxative should betaken with the first dose. Never put wafer into the mouth dry.	No. 1	15.1	5.6	None.	None.	47590	Acetanilide declared, but present in excess of amount named. Meets requirements of P. or P. Act.	
	" 2	15.1	5.6	"	"			
Directions:—Soften the wafer by placing it for a moment in a little water, then swallow with a drink of water. If relief is not obtained, a second wafer may be taken half an hour or an hour later.	No. 1	10.1	1.8	None.	None.	47591	Curative powers claimed. Meets requirements of the P. or P. Act.	
	" 2	10.0	1.8	"	"			