The subfamily <i>Ephutinæ</i> is divided into two tribes as follows:  Table of Tribes.	
Eyes never rounded or hemispherical, but always ovate, obovellipsoidal, not polished, and distinctly focusted	
Eyes rounded or hemispherical, very prominent and highly-po- not facetted, or the facets very indistinctly defined, as in the Photopsidini	hutini. lish <b>ed,</b>
Tribe I.—Ephutini.	
Table of Consu	
Males	
Females	9.
Eyes distinctly emarginate within.  Eyes not emarginate within.	
Eyes not emarginate within  2. Metathorax with the hind angles normal, not dentate; scut normal, not spined	
Metathorax with the hind angles dentate, clothed with a dense si white pubescence; scutellum bispined. (North and standard)	lvery-
3. Second and third cubital cells each receiving a recovery	Cam.)
transverse, or not longer than thick; first segment of abdopetioliform, as wide at base as at a con-	oints omen
Ephuta,	Sav.
= Rhoptromutilla, Ar	ndré.
Scape (?) not bicarinate beneath; first segment of the abdo	say.)
militario anterioriy, nodiform posterioris	
(Africa.)	ré.*
(Type R. clavicornis, And Second cubital cell receiving both recurrent nervures; so normal	iré.)
(Type Musilla	
Front wings with <i>three</i> cubital cells, or the third partially formed. Front wings with <i>two</i> cubital cells, the third entirely absent	

<sup>\*</sup>I have not seen a specimen of this genus, and am not positive of its position in this tribe.