Specimens collected in 1894 by Mr. A. P. Low at Lake Michikamow, Labrador, and in 1896 along the Koaksoak River, Northern Labrador, have been doubtfully referred here.

SAGINA NIVALIS, Lightf.

Along the Ungava River, Northern Labrador, 1896. (A. P. Low.) New to Canada.

HYPERICUM PROLIFICUM, L.

Sandy plains, along fence rows and in woods Caradoc township south of Strathroy, Ont., Aug., 1888. (J. Dearness.) New to Canada.

HYPERICUM NUDICAULE, Walt.:

H. Sarothra, Mx.; Macoun, Cat. Can. Plants, Vol. I., p. 85.Sandy fields south of Sandwich, Ont., 1892. (John Macoun.)Only authentic Canadian record.

GERANIUM BICKNELLII, N. L. Britton, Bull. Torr. Bot. Club, Vol. 24, p. 92.

"Similar to G. Carolinianum, but taller, the stems usually more slender, loosely pubescent. Leaves slender-petioled, somewhat angulate in outline, the segments oblong or linear-oblong, mostly narrower; peduncles slender, two-flowered, the inflorescence loose; sepals lanceolate, awn-pointed; ovary lobes pubescent; persistent filaments longer than the carpels; beak about 1' line long, long-pointed, its tip 2"-3" long; seeds reticulated."

"Nova Scotia (?) Maine to Western Ontario and Southern New York."

In the February number of the Bulletin of the Torrey Botanical Club, Dr. Britton describes G. Bicknellii as one of "two undescribed eastern species." Our herbarium sheets of G. Carolinianum have long been separated