

THE Canadian Journal of Medical Science.

A MONTHLY JOURNAL OF BRITISH AND FOREIGN MEDICAL SCIENCE, CRITICISM, AND NEWS.

U. OGDEN, M.D.,
EDITOR.

R. ZIMMERMAN, M.D., L.R.C.P., London,
107 Church Street, Toronto, Corresponding Editor.

SUBSCRIPTION, \$3 PER ANNUM.

All Communications, Letters and Exchanges must be addressed to the Corresponding Editor.

TORONTO. AUGUST, 1877.

Selections: Medicine.

THE MUSCULAR ARTERIOLES: THEIR STRUCTURE AND FUNCTION IN HEALTH AND IN CERTAIN MORBID STATES.

BY GEORGE JOHNSON, M.D., F.R.S.

LECTURE III.—*Concluded.*

There is good reason to believe that some of the more formidable *nervous symptoms*, which result from uræmia—in particular uræmic convulsions and a form of transient amaurosis—are directly due to cerebral anæmia consequent on sudden extreme contraction of the muscular arterioles.

There can, of course, be no question that uræmic convulsions are of an epileptic character. A large amount of evidence points to the conclusion that both the loss of consciousness and the convulsions of epilepsy are the results of sudden and extreme anæmia of the brain. In man, and in most, if not in all, warm-blooded animals, a rapid and very copious hæmorrhage usually causes convulsions. Kussmaul and Tenner state (*On the Nature and Origin of Epileptiform Convulsions caused by Profuse Bleeding*, New Sydenham Society, 1850) that in numerous cases of dogs, cats, and rabbits, they observed, without a single exception, violent and general convulsions preceding death by loss of blood. In order to produce this result, the hæmorrhage must be rapid. If it occur slowly, so that the vital powers are gradually exhausted, death then occurs with swooning, drowsiness, and delirium without convulsions.

The same experimenters found that an interruption of the supply of blood to the head of a rabbit, by ligature or compression of the arteries of the neck, produces epileptic convulsions as surely as hæmorrhage does. In about one hundred rabbits they ligatured or compressed the carotids and subclavians, from which, be it remembered, the vertebral arteries proceed; and in every case, except that of one very old animal lean and feeble, convulsions occurred.

In order to excite convulsions, they found it necessary to close all the four arteries which supply the brain. If but one carotid or one vertebral artery remained pervious, the animal was enfeebled and more or less paralysed, but not convulsed. And again, if, during the height of the convulsion, the ligature were removed from one carotid, the convulsions generally ceased immediately, and there was a sudden change from the most frightful spasm to complete relaxation of the muscles. The description of the convulsions thus artificially produced with, as it seems to me, needless reiteration, in the lower animals, shows that they were essentially the same as epileptic convulsions in the human subject. There was the dilated pupil, the tonic spasm, quickly followed by clonic convulsions so violent as to throw the animal forward to a distance of one or two feet, and sometimes even over the shoulders of the operator. The experiments obviously could not be performed on the human subject; but Drs. Kussmaul and Tenner approached as near to this as they dared by compressing the carotids of six men. The result was that in all the face turned pale; the pupils first contracted and then dilated; the respiration became slow, deep, and sighing; then there was giddiness,