

## 22.—RUFIOUS-CHEEKED WAGTAIL.

**SIZE** of the White Wagtail; length six inches and a half. Bill rather more stout than in that species, and dusky, with the base somewhat broad; plumage in general above brown; sides of the head under the eye, pale rufous; on the chin and throat a slight tinge of the same; the rest of the under parts white, passing on each side of the neck, but not quite to the back part; wings as the back, the four first quills nearly equal in length, but the first somewhat shorter; tail three inches long, much rounded, or slightly cuneiform; the feathers rather pointed at the ends; the outer one about half an inch shorter; colour rather darker than the wings, which, when closed, reach about one-fourth of the length; legs somewhat strong, one inch long, the joint fully covered with feathers, colour brown, claws crooked, moderate in size.

From the collection of Lord Stanley. Native place uncertain.

## 23.—GREEN WAGTAIL.

*Motacilla viridis*, *Ind. Orn.* ii. 505. *Gm. Lin.* i. 962.

Green Wagtail, *Gen. Syn.* iv. 403. *Brown, Ill.* 86. pl. 33. *Shaw's Zool.* x. 551.

**LENGTH** four inches. Head cinereous; neck, back, and breast, pale green; wings and tail cinereous, edged with white; belly white. Inhabits Ceylon.

## 24.—HUDSONIAN WAGTAIL.

*Motacilla Hudsonica*, *Ind. Orn.* ii. 503. *Vieill. Am.* ii. p. 47.

Hudsonian Wagtail, *Gen. Syn. Sup.* ii. 231. *Shaw's Zool.* x. 552.

**LENGTH** six inches. Bill somewhat stout, yellowish brown; plumage above brown, the feathers margined with ferruginous; chin and throat pale ferruginous, streaked with dusky; breast, belly, and