Where Joes it J Begin:

Is not a simple eastric derangement the precursor of most cases of Summer Diarrhoea in children? Are not the majority of these cases Dyspeptic in origin? Why not administer

<u>factobeptine</u>

early? you may "nip it in the bud." By righting the digestion and supplying the lacking ferments du you not pave the way for subsequent antiseptic treatment?

Can any course be more rational?

Has any other line of treatment proved more effectual?

Samples and literature upon request:



Hemaboloids
Blood Enricher

The lines of logic

INDICATE THE VALUE OF

Liquid Peptonoids

AS A SUITABLE SUBSTITUTE FOR FERMENTABLE MILK FOODS WHEN IT BECOMES NECESSARY TO DISCONTINUE THE LATTER IN CASES OF CHOLERIFORM DIARRHOEA ETC.

Liquid Peptonoids
IS THOROUGHLY PRE-DIGESTED
AND ITS QUICK ABSORPTION
IS THUS ASSURED.

ITS THOROUGHLY STERILE STATE PREVENTS ITS SEPTIC FERMENTATION.

ITS SLIGHT STIMULATING ACTION COMBATS COLLAPSE.

Liquid Peptonoids
IS THOROUGHLY PALATABLE AND TOLERABLE.

WHEN A COMBINATION OF A SEPTIC FOOD AND ANTISEPTIC PENEDRICS NEEDED

Liquid Peploncids

WITH CREOSOTE
WILL ME FOUND OF
PARAMOUNT VALUE

DEMOTOR CAMPLES

ARLINGTON CHEMICAL CONKER