

A summer resident, but not nearly so abundant as the last. Breeds in the interior of the Province.

A very destructive hawk.

SUBGENUS ASTUR.

135. **Goshawk.**

(*Accipiter atricapillus*.)

A winter visitor to Southern Ontario. Probably resident in the northern part of the Province.

This hawk is capable of killing full grown fowls and does so at every opportunity.

GENUS BUTEO.

136. **Red-tailed Hawk.**

(*Buteo borealis*.)

A common summer resident, breeding in the interior. Arrives early in March, departs at the beginning of November. A few have been seen in winter.

A useful hawk feeding chiefly on mice, grasshoppers and frogs.

137. **Red-shouldered Hawk.**

(*Buteo lineatus*.)

Common summer resident. Less frequently seen than the last. Breeds in the interior. Arrives a little later than the Red-tail, and leaves towards the end of October.

A useful hawk feeds chiefly on mice, reptiles and grasshoppers.

SUBGENUS TACHYTRIORCHIS.

138. **Swainson's Hawk.**

(*Buteo swainsoni*.)

A rare visitor from the western prairies where it is abundant.

139. **Broad-winged Hawk.**

(*Buteo latissimus*.)

A common summer resident. Breeds through its range. A useful species.

GENUS ARCHIBUTEO.

140. **American Rough-legged Hawk.**

(*Archibuteo lagopus sancti-johannis*.)