with a slight elevation. Wings greyish hyaline, due to the pubescence, venation brown, the 1st branch of the radius more than half as long as the 2nd, the recurrent nervure not interstitial with the 1st transverse cubital.

Described from one specimen.

(3) Bracon nigridorsum n. sp.

Q.—Length $3\frac{1}{5}$ mm. ; ovip. $\frac{1}{5}$ mm. Black, polished; mandibles, palpi, legs, including coxæ, and the abdomen, except the disk of 1st, 2nd, 3rd, 4th and 5th segments, which are black, wholly pale brownish-yellow. Tegulæ honey-yellow. Antennæ 35-jointed. Parapsides distinct. Metathorax and mesopleuræ smooth, polished, the latter with a single fovea near the posterior suture. Metapleuræ very hairy. Abdomen long oval, the shield of the 1st segment trapezoidal, finely rugose, rounded off at apex and between it and the lateral margins of the segment are two long channels; the 2nd segment is slightly shagreened at base. Wings hyaline, strongly iridescent, the venation brown; the 2nd branch of the radius is twice the length of the 1st, the recurrent nervure not interstitial.

SPATHIUS NEES.

(4) Spathius Canadensis n. sp.

 \mathcal{Q} .—Length 2_5^4 mm.; ovip. r_6^2 mm. Reddish-brown, the dorsum of thorax blackish or dark fuscous, the abdomen, except the long petiole and the base of the 2nd segment, black. The antennæ are honey-yellow, very long, multiarticulate, the posterior femora and tibiæ slightly fuscous. Head perfectly smooth, polished; the thorax delicately shagreened, the parapsidal grooves very distinct, deep; the metathorax with 3 or 4 delicate longitudinal carinæ. Wings fuscous, the base and tips hyaline and with a white or hyaline band across the middle, including the basal half of the stigma.

Described from one specimen. The species comes nearest to L. Laflammei Prov., but is readily separated from it and other species by its smaller size, perfectly smooth head and by the brevity of the ovipositor.

CÆNOPHANES FÖRSTER.

(5) Canophanes borealis n. sp.

 \mathcal{Q} .—Length 3_5^3 mm.; ovip. $\frac{4}{5}$ mm. Very elongate, black, finely rugose; the quadrate head is smooth, but with delicate transverse aciculations on