

Palpi	Slender. Nearly or quite twice long as eye.	Slender, compressed. Scarcely more than half as long again as eye.	Slender. Scarcely more than half as long again as eye.	Slender. Less than twice as long as eye.
(as the Nos.	2—	$1\frac{1}{2} +$	$1\frac{1}{2} +$	2—)
Ratio of fore tibia to hind	A little more than $\frac{5}{8}$.	$\frac{2}{3}$	A little more than $\frac{2}{3}$.	$\frac{3}{4}$
(as the Nos.	$.6\frac{1}{4}$	$.6\frac{2}{3}$	.6 +	$.7\frac{1}{2}$)
Ratio of mid tibia to hind	Scarcely $\frac{4}{5}$	$\frac{4}{5}$	Nearly $\frac{4}{5}$	A little shorter.
(as the Nos.	.8—	$.8\frac{1}{3}$	$.8\frac{1}{3}$ —	.9 or 1.0)
1st Sup'r Br. of subcostal ner- vure of fore wings arising	In the middle of the outer two-thirds of the upper border of the cell.	Somewhat beyond mid- dle of upper border of the cell.	At scarcely two-thirds the distance from the base to the apex of cell.	A little beyond the mid- dle of the upper bor- der of the cell.
2nd do.	Midway between it and the origin of 1st inferior subcostal nervure.	Midway between this & the origin of first in- ferior branch.	At less than half way between.	About one-fourth the distance from origin of first to apex of cell.
Cell	Somewhat more than $\frac{1}{2}$ as long as wing.	Somewhat more than $\frac{1}{2}$ as long as wing.	$\frac{1}{2}$ as long as wing.	Scarcely $\frac{1}{2}$ as long as wing.