

BAYER'S PHARMAGEUTICAL PRODUCTS

A tasteless, odourless, nutrient meat powder; it contains all the albuminoid principles of the meat in an easily soluble form. It has been extensively employed and found to be of the greatest service in consumption, diseases of the stomach and intestinal tract, chlorosis and rickets. It is of great value in convalescence from all diseases. SOMATOSE strengthens the muscles and stimulates the appetite in a remarkable manner. Dose for adults: a level teaspoonful three to four times a day with milk, gruel, coffee, etc.

IRON SOMATOSE (Ferro-Somatose).

A first-class tonic, containing the albuminous substances of the meat (albumoses) organically combined with iron. Special indications: Chlorosis and Anaemia. Daily dose: 75 to 150 grains.

MILK-SOMATOSE (Lacto-Somatose).

A strength giving food containing the albuminous matter (albumoses) of the

TRIONAL A most reliable and quickly-acting hypnotic of the Sulfonal group. Dose: 16 to 20 grains, in a large cup of hot liquid.

IODOTHYRINE The active principle of the thyroid gland. It is most efficacious in Strumous Diseases, Myxoedema, Obesity, Rickets, Psoriasis, Eczema, and Uterine Haemorrhages. Dose: 5 grains two to eight times a day for adults; 5 grains one to three times daily for children.

LYCETOL Tartrate of Piperazine

PHENACETINE-BAYER

PIPERAZINE-BAYER

Solvent. Has a marked effect on the diuresis. Dose: 16 to 32 grains daily.

ARISTOL An Iodine Cicatrisant which is an excellent odourless substitute for Iodoform and highly recommended for Burns, Wounds, Scrofulous Ulcerations, etc.

EUROPHEN A perfect substitute Odourless and non-toxic. Has a covering power five times greater than Iodoform. Especially useful in Ulcus molle et durum.

LOSOPHAN Α cresoltriiodide particularly efficacious in the treatment of all kinds of cutaneous disorders caused by animal parasites.

PROTARGOL A new silver preparation. Most reliable in cases of Gonorrhoea. Antiseptic wound healer. Excellent results in cases of Gonorrhoeal Ophtalmia. Solutions of ¼ to 2%. Ointments.

TANNIGEN An almost tasteless intestinal astringent. Most efficacious in Chronic, Acute and Summer Diarrhoeas. Adult dose: 8 grains every three hours.

TANNOPINE "Tan-(Formerly none"). intestinal astringent. Special indications: Tuberculous and non-tuberculous Enteritis, Typhus. Dose: 15 grains, three or four times daily.

SALOPHEN Specific for Influenza, Headache, Migraine, Acute Articular Rheumatism, Chorea, Sciatica. Dose: 15 grains four to six Anti-Arthritic, Uric times daily. In powders, etc.

> SULFONAL-BAYER SALOL-BAYER

Samples and literature may be had on application to the

DOMINION DYEWOOD CHEMICAL CO., TORONTO.

Sole Agency and Depot in Canada for all "BAYER'S" Pharmaceutical Products.

(WHOLESALE ONLY)