

Prothorax a very little deeper than the vertex ; hind margin concave, almost angled at the centre ; carinae very distinct ; colour uniform shiny reddish brown.

Legs brown, becoming lighter toward the distal end of the tibiae ; the tarsi yellow ; spur wide at base and very finely serrate.

Scutellum triangular, about twice as deep as the prothorax, sides arcuate, apex a rather sharp point, carinae distinct, the lateral carinae attaining the posterior margin, the median carina is obsolete toward the apex ; colour shiny, reddish brown, with the margins and apex yellow.

Elytra nearly twice as long as the abdomen ; colour smoky brown, veins darker brown with dark dots along them.

Tergum of the first two abdominal segments yellow, remainder of the abdomen reddish brown. Genitalia, aperture of the pygofer large, oval, wider than long ; ventral edge of the pygofer deeply notched, styles large, broad, divergent from their base, following the curve of the pygofer to the anal tube, a few long setae at their apical end. Length, including elytra, 3.5 mm.

Taken on reeds by the author in Renwick Swamp, Ithaca, N. Y., July 20, 1908. Five specimens are before me, one of these shows much more yellow on the vertex and front. Type in the author's collection.

This species suggests the macropterous form of *L. lineatipes* Van Duzee, but the colouring is different and, besides other minor differences, the genitalia are decidedly unlike that species.

*Achorotile foveata*, n. sp.

Macropterous ♀.—Eyes oval, deeply emarginate below to receive the antennae ; colour gray. Vertex as wide as long, rounded before, extending a little beyond the eyes ; carinae distinct, except the posterior median, which is weak ; posterior foveae deep ; colour yellowish brown, lighter posteriorly.

Front twice as long as wide, widest near the middle, sides gently curving ; clypeo-frontal suture straight ; the median carinae curved, following the curve of the sides to the front ; just outside of each median carina there is a row of pustules, six on each side, three near the vertex and three near the clypeus ; between these two groups are two pustules on each side along the outer margin of the front ; colour deep reddish brown.

Clypeus shiny black with an indistinct carina. Second segment of the antennae about one and one-third times as long as the first, covered with pustules ; basal segment reddish brown, second segment yellowish brown.