

rounded; subgenital plate asymmetrical, diagonally incised apically and furnished with short style-like organs; cerci short and stout, widest beyond the middle. The legs are black with a pre-apical yellowish band on the tibiae and the base of the basal segment of the tarsi also lighter.

Measurements. Total length from front of pronotum to the tip of the elytra, 7 mm.; of elytra 5.5 mm.; of wing, 10 mm.

Type a single ♂, Huascaray, Peru, September 21, 1911, altitude 6500 feet. C. H. T. Townsend collector. Catalogue No 15321 U. S. Nat. Museum.

*Cocconotus charape*, n. sp.

Description.—♂ (the ♀ unknown): Allied to *C. pulcher* Brunn. and runs to that species in the tables in Brunner's Monograph of the Pseudophilinae. It differs, however, very distinctly from that species.

In size and general coloration agreeing fairly well with *pulcher*. Head black and yellowish, the occiput blackish shading into yellowish on the cheeks and continuing yellowish down to the mouth-parts; mandibles, labrum and base of the clypeus and the sides of the face piceous, front of the face dark reddish brown with an apically pitted tubercle in the centre; on each side and just above the ends of the clypeal suture the face bears a large erect piceous pointed tubercle about as long as the clypeus; antennae piceous basally, shading gradually to reddish brown. Pronotum without carinae, the shoulders only slightly squared; disk slightly rugose, truncate behind, gently rounded before, the main transverse sulcus profound and situated distinctly behind the middle; prosternal spines long, sharp and piceous, the rest of the lower surface of the thorax light yellowish; the disk and the lateral lobes of the pronotum margined with piceous and the central portion of the disk, especially anterior of the principal sulcus, light yellowish brown, which colour continues down diagonally forwards entirely across the lateral lobes. Legs stout and yellowish, the coxae, the geniculations and the dorsal surface of the anterior tibiae more or less infuscated; fore tibiae furnished with conchate foramina and armed above on the inner margin with four tubercular swellings and armed beneath with a double row of spines; fore femora less than one and one half times as long as the pronotum, smooth above but armed beneath on the front margin with three short black spines; middle legs similar to the front ones but the tibiae have three distinct spinules above; hind femora very stout and short, the greatest width about three and one half times the length, smooth above, beneath armed on the outer side with seven or eight stout spines and on the inner margin with a smaller number of smaller spines, all the spines piceous to the base; hind tibiae