

the eyes a narrow line to the side of the neck, where it widens considerably, and continues in a broad patch to the back; this mark is at first black, but half an inch beyond the eye deep chestnut; chin and sides of the head, between the above line, white; neck dingy white, slightly tinged with chestnut; belly and all beneath white; at the back of the neck, between the two dark markings, a white line; back and scapulars dark ash, with a little mixture of chestnut; wings dark ash, larger coverts and secondaries slightly edged with white; the two middle tail feathers ash-colour, the others the same on the outer web; mottled ash-colour and white on the inner; upper tail coverts ash-colour, under white; legs black, bare near an inch above the knee; toes lobed, the outer united to the middle one, at a short distance from the base, claws small, curved, black.

Inhabits North America. Received in a collection dispatched from Cumberland House, in the spring, 1820.

---

**\*\* WITH FLAT BILLS.**

---

**3.—GREY PHALAROPE.**

- Phalaropus lobatus*, *Ind. Orn.* ii. 776. *Amer. Orn.* ix. 92. pl. 73. f. 3.  
*Tringa lobata*, *Lin.* i. 249. *Gm. Lin.* i. 674. *Muller*, No. 195.  
*Phalaropus*, *Bris.* vi. p. 12. *Id.* 8vo. ii. 361. *Fn. suec.* No. 179.  
*Phalaropus platyrhynchos*, *Tem. Man.* 459. *Id. Ed.* 2d. 712. *Parry's App.* p. cci.  
*Tringa cinerea gutture albo*, White-throated Coot-footed Tringa, *Bartr. Trav.* p. 302.  
*Le Phalarope à festons dentelés*, *Buf.* viii. 226.  
*Grey Coot-footed Tringa*, *Edw.* pl. 308. *Phil. Trans. L.* p. 255. pl. 6.  
*Grey Phalarope*, *Gen. Syn.* v. 272. *Br. Zool.* No. 218. pl. 76. *Id. fol.* 126. pl. E.  
1. f. 3. *Id.* 1812. ii. p. 123. pl. 21. *Arct. Zool.* ii. No. 218. *Bewick*, ii. pl. p.  
140. *Lewin*, v. pl. 194. *Orn. Dict.*

**LENGTH** seven inches and a half, extent sixteen; weight one ounce and three quarters. **Bill** black, flattened near the point, about