

10. *Ae. Nevadensis*.—A series of eight males from Reno, Nevada, are the only specimens seen.

The characters by which *Ae. interna*, *lineata* and *Nevadensis* are separated are so slight that it is by no means improbable that intermediate forms may be found. The first two named forms, however, are certainly characteristic of different geographical provinces, and although a considerable series of both has been examined, no intergrades have been seen. There seems to be no constant character for the separation of the females.

11. *Ae. eremita* ("Ae. X.," Williamson).—A very distinct species belonging to the wooded parts of the Boreal Zone from Labrador, Newfoundland and the White Mountains to the Hudson's Bay Slope, Great Slave Lake and Alaska, thence southward in the Rocky Mountains to Wyoming.

It occasionally wanders into the Transition Zone.

12. *Ae. clepsydra*.—Upper Austral and Transition Zones, from New England to Ontario and Indiana.

13. *Ae. Canadensis* ("Ae. Y.," Williamson).—Next to *Ae. umbrosa* this is the commonest *Aeshna* in collections, and is generally known as *Ae. clepsydra*. It is an abundant species in the Canadian division of the Boreal Zone, and is also common in the Transition Zone, from New England and the Maritime Provinces to Manitoba. Typical examples from Washington Territory have also been examined, and its range is therefore probably continuous from Atlantic to Pacific.

14. *Ae. verticalis*.—Very closely allied to *Ae. Canadensis*, but undoubtedly distinct. It inhabits the Transition and Upper Austral Zones east of the Mississippi.

15. *Ae. tuberculifera*.—A somewhat rare species, very distinct from any of the others. It has been taken in some numbers in the New England States, especially in southern Maine, and a few specimens from Ontario and Wisconsin have also been examined.

16. *Ae. juncea*.—A circumpolar species found in America in the subarctic forest belt, from Newfoundland to Alaska and south to the White Mountains, northern Ontario and in the Rocky Mountains to southern Colorado.

17. *Ae. subarctica*.—Boreal Zone, from Anticosti and Nova Scotia to Isle Royale, Mich., and the north shore of Lake Superior.

18. *Ae. sitchensis*.—Boreal Zone, from Newfoundland, through Quebec, northern Ontario, northern Michigan and Minnesota to Alaska.

19. *Ae. septentrionalis*.—Hudsonian division of the Boreal Zone, from Labrador, Newfoundland and the White Mountains to Great Slave Lake.