periodical and published in various journals, evince the estimation in which his talents are held by the reading public.

"Dr. John Boardman, of this city, takes Dr. Hunt's former position in the University as Demonstrator of Anatomy."

New Licentiates .- His Excellency the Governor General has been pleased to grant licenses to Charles Tozer, of Aylmer, in the county of Elgin, gentleman, and Thomas Benson, of Kingston, in the county of Frontenae, gentleman, to practise Physic, Surgery, and Midwifery in that part of Canada called Upper Canada.

Reviews .- Hereafter, we purpose placing the price of each be & we review immediately after the title. We do this for the benefit of our country subscribers, many of whom, we have been given to understand, would like to send for books if they were certain of the price. Mr. Dawson will transmit at y book ordered from him, through the post office. The postage is only one half-penny per ounce.

Books Received for Review .- Skoda on Auscultation and Percussion, 1854. From Messrs. Lindsay & Blakiston, Philadelphia. Annual Report of the City Inspector of the City of New York, for the year 1853. From T. K. Downing, Esq., City Inspector.

HOSPITAL REPORTS.

MONTREAL GENERAL HOSPITAL.

Case of advanced Iritis, rapidly yielding to treatment. (Reported by Mr W. J. Henry.

Mary Ann Gardner, a stout Irish woman, aged 30, of scrofulous appearance, married, was admitted into the Montreal General Hospital, on the 1st August, by Dr. Wright.

From the patient's statement, it appears that she caught cold on board a steamboat coming down from Hamilton about a fortnight ago, and a few days after her arrival in Montreal, felt a severe pain of a throbbing character, in one particular spot, just above the left eyebrow; this next passed to the temple-finally, the left eve became affected, and her sight began to be imperfect. She applied to a medical man here, and received some medicine from him, with directions to blister her temple. Experiencing no relief, she applied for admission into Hospital.

On examining the affected eye, a fine, light pink halo of vessels is seen radiating from the margin of the cornea; the iris is redder and