its lowto a conerest not en of the olete this articular

thin the red and le series, Trinity s detectof Come-

sections Random mination made of

sils were etion inestones, cheminards the

d upper p, while to the tter, he

vast acound, it be fully

he follination degrees

	FT.	IN.
a. Red and greenish slates chiefly dark-red with num-		
erous thin irregular calcareous layers	182	0
b. Band of greenish-gray slate	25	0
c. Red Slate	17	0
d. Bed of nodular-red and flesh-coloured limestone		
with shaley divisions, obscure fossils	4	0
e. Red slate	13	0
f. Thick bed light-red limestone	5	0
g. Red and green slates, chiefly red, a few thin layers		
limestone	38	0
h. Band of hard, greenish-gray sandstone	3	0
i. Wide band greenish shale	46	6
j. Red and green shale more red than green	53	6
k. Dark-red calcareous band weathering blackish very		
heavy, apparently containing a good deal of		
iron	2	0
l. Green. I and green shale in about equal propor-		
tions	57	0
m. Thick bed of hard, gray sandstone; may be a re-		
petition of band h	8	0
n. Red and green, shaloy slate alternating	66	0
o. Greenish-gray slates or shales containing fossils	145	0
p. Band Sesh-red limestone	1	0
q. Greenish-grey shale, darker towards top	<b>30</b>	0
r. Blackish, calcareous band, with peculiar, disk-like		
markings	10	0
s. Grenish-gray shales with thin, arenaeeous layers;		
numerous fossils	[୫୦	0
t. Black, finely laminated shales, with irregular, are-		
naceous layers much broken and contorted.		
Numerous fossils 4	25	9.0
1,2	51	0

The fossil organisms from this section and from the shales on Random Island having been sent to Mr. G. F. Matthew, of New Brunswick, for identification, have all been named and returned to the Museum. I have elsewhere referred to Mr. Matthew's uniform