ABRONIA LATIFOLIA. Esch.

Found by many collectors in the vicinity of Victoria, Vancouver Island. Very fine specimens of this species were collected on the west coast of Vancouver Island at Long Beach, north of Ucluelet by Mr. Wm. Spreadborough in 1909. Roots were found which in their dried state are three inches in diameter, they were nearly six inches in diameter when collected.

BIRD MIGRATION, 1908. OBSERVATIONS MADE ON SABLE ISLAND, NOVA SCOTIA. By R. S. BOUTEILLIER.

	When first		Number
	see		seen.
Semipalmated Plover	. April	17	2 or 3.
Robin	. "	20	about 12.
White-throated Sparrow	. "	22	several.
Tern		25	several.
Piping Plover	. May	10	2.
Least Sandpiper	. "	14	2.
Roseate Tern	. "	14	2.
Swallow		14	a few.
Martin		14	a few.
House Sparrow		21	2.
Phalerope	**	22	about 20.
Black-throated Green Warbler		25	1.
Black-polled Warbler		25	1.
Maryland Yellow-throat		25	1.
Blackbird	Tune	2	1.
Greater Yellowleg	Inde	8	2.
Pectoral Sandpiper	July	22	1.
White-rumped Sandpiper		24	30.
Black-bellied Plover	Δ	16	3.
Golden Plover	.Aug.	20	
Yellow-rumped Warbler			several.
Black-throated Blue Warbler		15	2 or 3.
		16	1.
Least Tern		16	several.
Blue-winged Teal		19	2.
Pine Grosbeak		19	1.
Sora Rail		19	1.
Pipet	. "	19	several.
Blackbird	. "	19	1.