

We omit purposely all references to Mr. Scudder's notices of these species. Two brief notices of the occurrence of *T. falacer* in New England preceded the paper in the Boston proceedings noticed, and to which the present is a reply.

With respect to the geographical distribution of the two species, while occurring side by side in the Atlantic District, it is probable that *Thecla inorata*, already found from Canada to Georgia, may be found over a wider expanse of territory than *Thecla falacer*.

Demopolis, Ala., December, 1870.

INSECTS OF THE NORTHERN PARTS OF BRITISH AMERICA.

COMPILED BY THE EDITOR.

From Kirby's Fauna Boreali-Americana: Insecta.

(Continued from page 145.)

67. TRECHUS [BRADYCELLUS] TIBIALIS, Kirby.—Length of body $2\frac{3}{4}$ lines. Only a single specimen taken.

[47] Body black, somewhat glossy. The tip of the palpi and scape of the antennæ are rufous; the prothorax is rather wider than long, but nearly square; the short basilar furrows observable in *Argutor* distinguish this species from the succeeding ones: elytra lightly furrowed, furrows impunctured; in the usual situation adjacent to the second furrow a little beyond the middle of the elytrum a very minute punctiform impression is just discernible; the lateral margin and suture at the apex of the elytra are reddish: the tibiæ are rufous but the cubit is black at the tip; the tarsi are darker, the hand has four dilated joints as in the other species of the genus.

[Belongs to *Bradycellus*; for a synopsis of the N. American species by Dr. LeConte, *vide* Pro. Acad. N. S. Phil., Dec. 1868, p. 379.]

68. TRECHUS [BRADYCELLUS] RUFICRUS, Kirby.—Length of body $2\frac{1}{2}$ lines. Only one specimen taken.

Body black, glossy. Palpi piceous; scape of the antennæ and mouth rufous: prothorax subobcordate; channelled, channel not abbreviated, margin rufous especially the basilar; angles rounded; basilar impressions single, round and punctured: elytra lurid or dirty yellow, with a large blackish cloud or blotch beyond the middle: legs dull rufous, thighs darker. [Included by LeConte, *loc. cit.*, under *B. badüpennis*, Hald., a species taken at Grimsby, Ont., by Mr. Pettit.]

69. TRECHUS [BRADYCELLUS] FLAVIPES, Kirby.—Length of body $2\frac{1}{2}$ lines. Many taken in Lat. 54° .

[48] Body piceous, glossy. Head underneath, nose, mouth, and oral organs—except the palpi which are pale yellow—and antennæ rufous; three