

Table 1

Infant Death rates by major cause, New Brunswick and Canada, 1963-1967,  
(per 1000 live births)

	1963		1964		1965		1966		1967	
	N.B.	Canada								
Total . . . . .	27.8	26.3	26.1	24.7	23.0	23.6	24.1	23.1	25.1	22.0
Lower respiratory infections . .	4.8	3.5	5.1	3.0	4.0	2.9	3.6	2.7	3.3	2.5
Immaturity . . . . .	6.6	5.0	7.2	4.8	5.4	4.4	6.0	4.2	6.7	3.9
Congenital malformations . . . .	5.7	4.4	4.2	4.4	3.5	4.3	5.1	4.5	3.2	4.0
Birth injury . . . . .	2.0	2.7	2.2	2.6	1.6	2.4	1.7	2.3	2.4	2.1
Asphyxia and atelectasis . . . .	1.9	2.9	1.8	2.7	1.8	2.6	2.1	2.5	1.4	2.4
Gastro-intestinal infections . .	0.6	1.0	0.5	0.6	0.6	0.5	0.5	0.5	0.7	0.5
Ill-defined . . . . .	2.4	2.5	1.4	2.4	1.8	2.5	1.7	2.6	2.9	2.8
Other . . . . .	3.8	4.3	3.7	4.2	4.3	4.0	3.4	3.8	4.5	3.8

SOURCE: Vital Statistics D.B.S. 84-202

Table 2

Perinatal Death\* Rates, New Brunswick and Canada  
1963 - 1967  
(per 1000 total births)

	1963	1964	1965	1966	1967
N.B. . . . .	30.1	29.4	25.7	26.8	28.4
Canada . . . . .	28.0	27.4	26.0	25.5	24.7

\* Foetal deaths (stillbirths) of 28+ weeks gestation and infant deaths under 1 week

SOURCE: Vital Statistics D.B.S. 84-202