

*Stibadium manti*, n. sp.—Expanse : 31 mm.

General colour walnut brown, quite dark over outer part of median space, and a triangular patch just before apex, paler basally and submarginally, though the contrast is not strongly pronounced. Under the lens there is seen to be a more or less evident sprinkling of white scales. Head, collar and thorax concolorous. Ordinary lines and spots not prominent, except t. a. and t. p. lines, and especially the latter, they are hard to make out except with lens. Basal half line not in evidence in any of the specimens. T. a. inwardly oblique, slightly waved or scalloped, subobsolete in some specimens, and more in evidence from the slight accumulation of pale scales along the outer side, than from the only slightly darker brown of the line itself. The t. p. line is better marked in most specimens, and in the proper light can be quite readily traced with the naked eye. Starting from costa, it makes a slightly downward curve to a point some little distance beyond cell, whence, after making a sharp, though rounded angle, it proceeds with a very slight inward curve in a strongly oblique direction to inner margin. The line itself is a trifle darker than the ground colour, but is emphasized by an outer border of white scales. In one quite fresh specimen the line can scarcely be traced, even with the lens, but the contrasting median and subterminal spaces mark its position. In one or two specimens very faint traces of a submarginal line, pale and irregular, can be made out with the lens. Orbicular very faint, concolorous or slightly darkened, round, outlined by white scales. Reniform subobsolete, the inner margin usually in evidence, the outer fragmentary or wanting. Claviform wanting. Fringe brown, paler opposite ends of veins. Secondaries quite uniformly dark fuscous brown, slightly darker along outer border, a very faint extra-mesial pale band can be made out in the proper light. Fringe fuscous, paler at base and slightly so at ends of veins. Beneath, pale grayish brown, more gray along costa and outer border, more brown centrally, faint extra-mesial band. Hind wings paler, quite gray, mesial band very faint.

Types : ♂ and ♀, Kerrville and San Antonio, Texas. From Mr. Lacey and Professor Attwater.

*Ogdoconta altura*, n. sp.—Expanse : 21 to 25 mm.

General colour of head, collar, thorax and fore wings a dark golden brown, thinly sprinkled with pale scales. Basal half line only apparent under lens, and then only as a few pale scales. T. a. line plainly visible but not prominent, pale, inwardly oblique, slightly outcurved. T. p. line the only contrasting feature of wing, pale, angled close to costa, then