with the collar white, the front and middle legs below the knees rufous infuscated. Length, 6.5 mm.

Habitat.-Olympia, Washington. Professor Trevor Kincaid, collector.

Empria concisa, n. sp.—Female. Clypeus with a median carina, roundly emarginate without a tooth, lobes rounded; the antennal furrows not interrupted; the ocellar basin broad, with sloping sides, reaching the median ocellus, extending to the middle of the antennal foveæ, the median fovea a small pit on a level the dorsal margin of the antennal foveæ; the ocellar and interocellar furrows distinct; the third segment of the antennae longer than the fourth; the saw-guides convex above and below, obliquely rounded at apex above and below to a point slightly above the middle of the apex; body black, with the labrum, collar, tegulæ, knees, front and middle tibiæ and tarsi, all more or less infuscated, white. Length, 7 mm.

Habitat.—Pullman, Washington. Professor C. V. Piper, collector.

No. 13.

Empria cupida, n. sp.—Female. Clypeus with a median carina, shallowly emarginate with a tooth, lobes acute; the ocellar basin a broad, shallow depression, broadest near the middle, extending to the median ocellus; median fovea broad and deep, situated above the dorsal margin of the antennal foveæ; ocellar and interocellar furrows distinct; the third segment of the antennæ longer than the fourth; the antennal furrows interrupted; the saw-guides straight above, convex below, obliquely, convexly rounded to a point above at apex; body black, with the collar and front and middle legs below the knees white. Length, 6 mm.

Habitat.—Olympia, Washington. Professor Trevor Kincaid, collector.

MISCELLANEOUS NOTES ON THE HYMENOPTERA CHAL-CIDOIDEA: THE GENUS ARTHROLYTUS THOMSON; HORISMENUS MICROGASTER ASHMEAD.

BY A. ARSENE GIRAULT, UNIVERSITY OF ILLINOIS.

Introduction.

In describing a new species of Arthrolytus Thomson, I merely bring together the literature of the group.

History and Description.

The genus Arthrolytus was founded by C. G. Thomson in 1878 with the rank of a subgenus Pteromalus Swederus, 1795. The description of Pteromalus and the Conspectus subgenerum, which includes Arthrolytus, were given by Thomson as follows:

October, 1911