

are often also useful, and a generous diet is always indicated. Unless the stomach has passed into a state of disease (which it may do, if overtaken when in this weakened state), any of these medicines are generally well borne. It will be well to bear in mind, however, that if the mucous membrane of the stomach be in a state of irritation, quinine, arsenic, phosphorus, the hypophosphites, and sometimes even cod-liver oil, are generally inadmissible."—*British Medical Journal*.

ON THE THERAPEUTIC VALUE OF SULPHUROUS ACID IN SCARLATINA MALIGNA.

Dr. Keith Norman Macdonald, after denying the prevalent opinion that no reliance can be placed on any drug in cases of scarlatina, does not hesitate in affirming that, when properly applied, both locally and internally, sulphurous acid is by far the most efficacious remedy we possess. He continues: "I have had several opportunities of testing its efficacy in some of the worst cases I have ever seen, during the epidemic which has been rife in this town (Cupar Fife) for the last two months, and I am bound to say that, of all remedial measures in this disease, it is, in my opinion, the most reliable. My treatment is as follows. The moment the throat begins to become affected, I administer to a child, say of about six years of age, ten minims of the sulphurous acid, with a small quantity of glycerine in water, every two hours, and I direct the sulphurous acid spray to be applied every three hours to the fauces for a few minutes at a time, by using the pure acid, in severe cases, or equal parts of the acid and water, according to the severity of the case. Sulphur should also be burned in the sick chamber half a dozen times a day, by placing flour of sulphur upon a red hot cinder, and diffusing the sulphurous acid vapour through the room, until the atmosphere begins to become unpleasant to breathe.

"In the worst cases, where medicine cannot be swallowed, this and the spray must be entirely relied upon; and the dark shades which collect upon the teeth and lips should be frequently laved with a solution of the liquor potass permanganatis, of the strength of about one drachm to six ounces of water, some of which should be swallowed, if possible.

"In cases presenting a diphtheritic character, the tincture of perchloride of iron should be administered in rather large doses in a separate mixture with chlorate of potash, and equal parts of the same with glycerine should be applied locally, with a camel's-hair brush several times in the day; but, as in the majority of cases among children, it is next to impossible to use a local application more than once; the spray and permanganate solution will then prove of great service.

"As to other remedies recommended by various authors, ammonia is nasty, and cannot be taken well by children; carbolic acid has the same fault, and cannot be applied properly. Gargles are also useless in children, because they seldom reach the diseased surfaces, and warm baths and wet sheet packing are dangerous, because they are never carried out properly in private practice. The hypodermic injection of pilocarpine is a remedy that may give good results hereafter, but I have had no experience of its use."—*British Medical Journal*.

DISLOCATIONS OF THE THIGH REDUCED BY NEW METHODS OF MANIPULATION.

In cases where reduction of the femur by manipulation, in the usual way, with the aid of anæsthetics, has failed, or is inapplicable, and as a substitute, in many cases, for anæsthesia, assistants, and mechanical power, Mr. Kely (*Dublin Journal of Medical Science*, October) proposes the following methods:

For posterior dislocations.—The patient is laid prostrate upon the floor. Three strong screw-hooks are inserted into the flooring close to the perineum and each ilium of the patient, and to these hooks he is secured by strong bandages or rope. The injured thigh is flexed at right angles to the patient's body: the foot and lower extremity of the tibia are placed against the perineum of the surgeon who, bending forward, with the knees slightly flexed passes his forearms behind the patient's knee and grasps his own elbows. Reduction is now accomplished by drawing the femur upwards; but circumduction may also be practised; the surgeon, stepping backward, then extends the limb, and lays it by the side of its fellow. In sciatic dislocations, in order to liberate the head of the bone from the foramen, a bandage may be passed around the thigh, close to the trochanter, by which an assistant may make traction.

For anterior dislocations.—The patient is placed upon a table of such elevation as to have his pelvis nearly as high as the trochanter of the surgeon. A bandage around the pelvis, and secured to the side of the table farthest from the dislocation, affords counter-extension. The surgeon, with his face directed towards the dislocated joint, and standing on its inner side, with his trochanter pressed against the femur, now bends the leg behind his back, and grasps the ankle with the corresponding hand. Reduction is effected by rotating or turning his body partially away from the patient, thus making traction on the femur in the most favorable direction, and at the same time pressing its head towards the acetabulum with the disengaged hand.