

*Vanessa antiopa*, Linn.

Very rare ; have seen but three specimens, one in March, one in July and one in November.

*Pyrameis atalanta*, Linn.

Very common from April to October.

*Pyrameis huntera*, Fabr.

The least common of its genus but not uncommon, and is found from May to October.

*Pyrameis cardui*, Linn.

Very common during the same months.

*Junonia cœnia*, Hbn.

Uncommon, but a few specimens seen every year from July to October.

*Limenitis ursula*, Fabr.

Not uncommon from May to October.

*Limenitis dissippus*, Gdt.

Found mostly in August around maple trees ; not uncommon.

*Apatura celtis*, Bd.-Lec.

Not common and is found in numbers only in a yard in town which has a hackberry tree. August is when most are seen.

*Debis portlandia*, Fabr.

Common all through the summer months.

*Neonympha eurýtris*, Fabr.

. Common from May to September, and is found chiefly near the bottom of gullies which have grassy sides.

*Satyrus alope*, Fabr., var. *Olympus*, Edw.

Very rare ; have seen but two specimens.

*Libythea bachmani*, Kirtl.

Usually not uncommon through the summer months.

*Thecla melinus*, Hbn., var. *Humuli*, Harr.

Very rare ; have taken but one specimen.

*Feniseca tarquinius*, Fabr.

Very rare ; took one specimen, somewhat broken, in an opening in the woods in September.

*Chrysophanus thoe*, Bd.-Lec.

Rare ; a few specimens were captured by a friend west of Marshall in September.