

TABLE B.
Genera of the Solanum-Cassia Type. Distribution.

Region.	Endemic.	Others.	Total.
1	2 = 3.38%	8	10 = 16.94%
2	1 = 1.69%	7	8 = 13.55%
3	7 = 11.86%	9	16 = 27.11%
4	17 = 28.81%	9	26 = 44.06%
5	2 = 3.38%	3	5 = 8.47%
6	12 = 20.33%	6	18 = 30.50%
7	4 = 6.77%	4	8 = 13.55%
8	2	2 = 3.38%
9
10	2	2 = 3.38%
11	2	2 = 3.38%
12	1 = 1.69%	4	5 = 8.47%
13	4	4 = 6.77%
14	1	1 = 1.69%

We find that the Indian region has 11.8 per cent., the South American 28.8 per cent., the Australian region 20.3 per cent., and the extra-tropical South American region 6.7 per cent. of the members of this type confined exclusively to their limits. The average for these four regions is 16.9 per cent., while the average for the other ten regions is .9 per cent., six of the regions having no endemic representatives of this type at all. Considering all representatives of the Solanum-Cassia type occurring in the several regions, whether endemic there or not, we find that the Indian region contains representatives of 27.1 per cent. of the 59 genera, the tropical South American region 44 per cent., the Australian region 30.5 per cent., and the extra-tropical South American 13.5 per cent. The average for