

and legs brownish yellow. Head very finely and sparsely punctate, not tuberculate; clypeus broadly and shallowly emarginate at middle, the sides curved. Thorax not narrowed in front, sides broadly curved, hind angles obtusely rounded; base very distinctly margined, disk finely and sparsely punctate. Elytra equal in width to thorax, finely striate, the striae finely and indistinctly punctate; intervals feebly convex, smooth. Front tibiae stout, broad, distinctly punctate in front, strongly 3-toothed. Hind tibiae rather slender; first joint of hind tarsi as long as the next three together. Length 3 mm.

Two specimens from beneath dry cow-dung in company with *A. vestiarius* Horn, near Sarasota, February 17th. Closely related to *A. rubeolus* Beauv., but smaller, more slender, with paler elytra, longer basal joint of hind tarsus, and with base of thorax distinctly margined. One of the types is in the collection of F. Blanchard, Tyngsboro, Massachusetts.

*Hymenorus granulatus*, sp. nov.

Oblong-parallel, subdepressed, sparsely pubescent with fine recumbent grayish hairs. Black, shining; palpi and mandibles reddish brown, tarsi piceous. Eyes large, separated by about one-half their own diameters; antennae stout, half the length of the body, the joints flattened, triangular, the third twice as long as second, half the length of fourth. Thorax at base one-third wider than long, sides broadly rounded into the front margin, base slightly sinuate each side near middle; disk strongly declivent in the region near the front angles, obsoletely foveate each side of middle at base, very densely and coarsely punctate, the punctures feebly separated or in part confluent. Elytra slightly wider than base of thorax, their sides parallel to apical fourth; disk striate, the intervals convex, densely granulate-punctate. Basal joint of hind tarsi slightly curved, one-half longer than the remaining joints together. Length 7—7.5 mm.

Described from four specimens beaten from scrub-oak foliage near Sanford, March 28—29. The dense punctuation of thorax and elytra, taken in connection with the uniform shining black color, readily distinguishes this from all other described species of *Hymenorus*.

*Andrimus confusus*, sp. nov.

Elongate-oval, sparsely clothed with short, suberect yellowish hairs. Head, thorax, under surface and legs reddish brown; elytra and antennae dark chestnut brown. Head transversely sulcate in front of eyes, finely and rather closely punctate, alutaceous between the punctures; eyes large,