



Fig. 2 The imperfectly formed urethra in a pseudohermaphrodite. The glans penis is perfectly formed in its upper portion and above is provided with a rudimentary prepuce formed by the labia. The penile portion of the urethra is recognized in part as a groove, in part as a ridge. The urethral orifice is of the female type and is freely patulous to the bladder. The vagina is $1\frac{1}{4}$ inches deep and ends in a blind pocket. There is no evidence of a cervix.

of spermatozoa. The stroma of the testicle was increased in amount.

The epididymis was little altered.

ANATOMICAL POINTS OF DIFFERENCE BETWEEN THE NORMALLY FORMED FEMALE AND OUR PATIENT

Mr. Brödel, who had just read Stratz's book¹ on the female human body, suggested that it might be of interest to compare the superficial anatomical landmarks in our case with those laid down by Stratz as constituting the normal in the female. The accompanying tabulations give the

interesting results as determined by Mr. Brödel after a most careful examination.

MEASUREMENTS OF AVERAGE-SIZED FEMALE (A) AND LARGE FEMALE (B) COMPARED WITH PROPORTIONS IN OUR PATIENT (C), THE BODY LENGTH OF B AND C CORRESPONDING.

	A	B	C
	Normal female of av. size	Normal female of size of our patient	Measurements of our patient
Length of body.....	168 cm.	183 cm.	183 cm.
Weight of body.....	132 lbs.	148 lbs.	170 lbs.
From nose to symphysis (length of spinal column),.....	62.5 cm.	68.5 cm.	71.5 cm.
Length of head.....	21 cm.	23 cm.	24 cm.
Width at temples.....	13 cm.

¹Stratz, C. H. Die Schönheit des Weiblichen Körpers. Stuttgart, 1900.