

Wings smoky golden, all except internal border; tegulæ golden fuscous, halteres concolorous.

Length of body, 5 mm.; of wing, $4\frac{1}{2}$ mm.

Described from one specimen; So. Illinois (Robertson).

Ennyomma distoides, Twms. Trans. Am. Ent. Soc., XVIII, p. 371.

A ♂ specimen from So. Dakota (Aldrich), August 12. Measures $7\frac{1}{2}$ mm.

Clista americana, n. sp., ♀.

Eyes brown; frontal vitta dark brown, blackish, averaging one-third width of front, front averaging one-third width of head; sides of front, sides face and facial depression black, thinly silvery-pollinose; cheeks posteriorly black, thinly silvery, rest included in the large brown pregenal area, an angle of which runs upward between sides of face and facial depression; sides of face fringed with bristles on inner border, cheeks on lower border; antennæ entirely rufous, third joint about twice as long as second, arista brownish; proboscis and palpi brown or blackish, the proboscis fleshy, not as long as height of head, the palpi curved and thickened at tip; occiput black, black-hairy. Thorax and scutellum shining black, the thorax in front slightly silvery, leaving three black vittæ which become lost near suture. Abdomen shining black, with a hardly greenish reflection, broad, flattened, rounded; first two segments with a lateral and a median marginal pair of weak macrochætæ; third with about eight marginal, and anal about six marginal macrochætæ of normal size. Legs black, claws and pulvilli only a little elongate, pulvilli smoky-yellowish. Wings grayish-hyaline, base and veins yellow; tegulæ brassy-yellow, front scales white on outside, the rest transparent; halteres yellow, tinged with rufous toward base.

Length of body, $6\frac{1}{2}$ mm.; of wing, $5\frac{1}{2}$ mm.

Described from one specimen; So. Illinois (Robertson).

Tryphera americana, n. sp., ♀.

Eyes bare, brownish; front more than one-third width of head, frontal vitta velvety brown; frontal bristles descending to base of third antennal joint, vertical bristles strongest, and with three next pairs directed backward, two orbital bristles; sides of front shining black; facial depression and sides of face black, somewhat silvery, epistoma pale luteous; cheeks shining black, anteriorly somewhat rufous; facial ridges bare; vibrissæ decussate, inserted on oral margin; antennæ rufous, more or less blackish, third joint blackish-brown in some lights, fully two and one half times as long as second, somewhat widened; arista brown, 3-jointed, often some-