

Sir B. Brodie and Mr. Gray, Med. Chirurg. Trans., vol. 39. }
Disease had extended into joint and attacked patella and tibia.

None for 10 years.

{ 10 years and not said to be dead, }
{ first felt. }

{ Condyles femur, patella and head tibia. }

M.

36

28

29	26	F.	Condyles femur.	Not stated.	Not stated.	Not stated : had it 5 years before removal?	None for 10 years.	May, '54.	Mr. Ward and Mr. Adams, Path. Trans. vol. v.	{ Healthy; except absorption of cartilage to extent of 3d. } Not stated.
30	28	M.	Condyles femur.	No injury.	Alive 10 years after.	Alive 14 years after.	None for 10 years.	1844.	Mr. P. Hewitt.— <i>Id.</i>	Not stated.
31	74	F.	D. matter over pectrous bone.	None apparent.	Not stated.	{ Died of pneumonia about 2 mos. after removal. }	Head symptoms.		Lebert; Phy. Path. t. 2, p. 148.	
32	46	F.	About great toe, not invol'g bone.	None apparent.	Not stated.	20 years before removal.	Not stated.		Do.	
33	28	F.	Periosteum, lower part leg.	None apparent.	Alive 7 yrs. after.	Alive 11 years after.	None for 7 years.	Dec., '38.	Do.	
34	12	F.	Head tibia.	Appeared 2 mos. after an injury.	Alive 6 mos. after.	Alive 3½ years.	None for 6 mos.	March, '56	Mr. Shaw; Patholog. Trans., vol. vii.	Not stated.
35	19	M.	Lower end fibula.	None apparent.	Alive 4 yrs. after.	Alive 5 yrs. after.	None for 4 years.		Messrs. Parker, Paget and Gray; Medical Chir. Trans., vol. 38.	Do.
36	40	F.	Attached to lower end radius.	Do.	Alive 2 mos. after.	Alive 12 to 15 years after.	None for 2 months after complete removal.	Dec., '51.	Mr. B. Cooper; Med. Times & Gaz., Feb., 1852, p. 214.	Healthy.
37	32	M.	Head fibula.	Do.	{ Alive 2½ years died few days after 2d removal (val of pleurisy } 10 days when reported.	2½ years after first removal, 3 myeloid tumors in stump.		Nov., '58.	Mr. Cock & Dr. Wilks; Med. Times & Gaz., Jan., 1859, p. 71.	Cartilage perfect.
38	17	F.	Condyles femur.	Frequent blows.	10 days when reported.	Alive 8 or 9 mos. after.		Jan., '59.	Mr. Simon and Sidney Jones.— <i>Id.</i> Feb., '59, p. 171.	{ Growth occupied joint, but cartilages intact }

17

CASES OF A SOMEWHAT DOUBTFUL NATURE.

39	53	M.	Neck beneath sterno-mastoid.	None known.	Not removed.	1 year.	{ Died exhausted; 4 small myeloid masses in lungs, and diffused in 1 cervical gland. }		Mr. Paget; Surg. Path., p. 454.	
40	33	F.	Head humerus.	Appeared 14 mos. After fracture and dislocation.	Died 5 mos. after.	5½ years.	{ 10 weeks after removal recurred as cancer at site of wound and in lungs. }		Mr. Hutchinson; Pat. Trans., vol. viii.	Not stated.
41	50	F.	Mamma.	Not stated.	Died 21 months after.	2½ years.	{ Glands enlarged, but not cancerous. }			
42		M.	Head fibula.	None known.	Died few m'ths after.	{ Operated on 6 mos. after discovery of tumor, and 2-12 yrs. after first occurrence of pain. }	6 months as open cancer of axilla.		Paget; Sur. Path.	
							{ A few months after operation myeloid and cancer found in lungs and spine. }	May, '57.	Mr. Cocks & Dr. Wilks; Guy's Hos. Reports, vol. iii., p. 175, and vol. iv., p. 31.	Not stated.