

Intervals perspiration broke out, chiefly on the forehead and chest, when the pupils were tending to dilate but disappeared when he became more fully under the influence of the Calabar bean. The urine was drawn off three times, the first time without causing spasm. Chloral was given (grs. xxx,) three times during the day. In the afternoon the bowels were freely opened; he was quite conscious of the motion. He became quite restless in the afternoon and evening; the pupils were constantly tending to dilate, and he was with difficulty kept under the influence of the bean; whenever the pupils dilated or even showed a tendency to dilate the spasms at once returned, and became rather more severe. During the twenty-four hours the temperature ranged between 97.8° and 100.2° ; the pulse between 98 and 130; and the respirations between 26 and 34. The spasms occurred at:

A.M.	A.M.	P.M.	P.M.	P.M.
12.30	9.30	12.35	4.15	7.35
12.45	9.45	12.45	4.45	8.05
1.30	10.45	1.00	4.50	8.20
3.15	10.58	1.15	4.55	8.55
3.30	11.10	2.00	5.00	9.50
4.30	11.29	2.08	5.15	10.20
4.50	11.45	2.20	5.30	10.30
5.10	11.50	2.30	5.45	10.40
6.10	12.00	3.05	6.00	10.55
7.30	P.M.	3.20	6.15	11.25
7.45	12.05	3.40	7.05	11.30
8.15	12.25	3.55	7.25	11.45

Up to midnight $42\frac{1}{2}$ grains had been given hypodermically.

Aug. 29th.—During the night the pupils manifested a tendency constantly to dilate; the bean was administered hypodermically, $\frac{1}{2}$ a grain every quarter of an hour as the condition of the patient seemed to require. After midnight, the spasms were slight though frequent; they in fact resembled cramps rather than true spasms. Chloral (gr. xxx) was administered twice during the night; the urine was drawn off once causing slight spasm; the bowels were freely opened twice. Although restless at times, he slept tolerably well. All forenoon and afternoon, the pupils kept dilating and contracting, and the