## Where Joes it Jegin:

Is not a simple Eastric derangement the precursor of most cases of Summer Diarrhoea in children? Are not the majority of these cases Dyspeptic in origin? Why not administer

lactopeptine

early? you may "nip it in the bud." By righting the disestion and supplying the lacking ferments du you not pave the way for subsequent antiseptic treatment?

Can any course be more rational?

Has any other line of treatment proved more effectual?

Samples and literature upon request:



Hemaboloids
Blood Enricher

## The lines of logic

Signid Peblonois

AS A SUITABLE SUBSTITUTE FOR FERMENTABLE MILK FOODS WHEN IT BECOMES NECESSARY TO DISCONTINUE THE LATTER IN CASES OF CHOLERIFORM DIARRHOEA ETC.

Liquid Peptonoids
IS THOROUGHLY PRE-DIGESTED
AND ITS QUICK ABSORPTION
IS THUS ASSURED

ITS THOROUGHLY STERILE STATE PREVENTS ITS SEPTIC FERMENTATION.

ITS SLIGHT STIMULATING ACTION COMBATS COLLAPSE.

Liquid Peptonoids
IS THOROUGHLY PALATABLE AND
TOLERABLE.

WHEN A COMBINATION OF A SEPTIC FOOD AND ANTI-SEPTIC REMEDY IS NEEDED

Liquid Peploncids

WITH CREOSOTE WILL BE FOUND OF PARAMOUNT VALUE.

SEND FOR SAMPLES.

THE ARLINGTON (HEMICAL