

on under the influence of air and water. Such is the case with the alkalies, potash and soda, and to a certain extent with the phosphates. Now although there is probably no soil which does not yield by analysis quantities of all the mineral elements sufficient for many crops, yet by long and uninterrupted tillage the more soluble combinations of these elements may be all taken up, and the land will then require a certain time of repose in order that a store of more soluble matters may be formed. Hence the utility of fallows.

In my analyses of the soils of the Richelieu valley, in the Report for 1850, pp. 79—90, I have shown, by comparing the virgin soils with those exhausted by continued crops of wheat during fifty years, that the proportions of phosphoric acid and magnesia, elements which are contained in large quantities in this grain, have been greatly diminished, but the soil still contains as much phosphate as it has lost, and this only requires to be rendered soluble in order to be available to vegetation.

In forests and untilled lands the conditions of a healthy vegetable growth are seldom wanting; the soil affords in sufficient quantity all the chemical elements required, while the leaves and seeds which annually fall and decay, give back to the earth a great proportion of the elements which it has yielded. In this way the only loss of mineral matter is that which remains stored up in the growing wood or is removed by waters from the soil. Far different is the case in cultivated fields, in the shape of corn, of fat cattle, and the products of the dairy, we remove from the soil its phosphates, alkalies and nitrogen, and send them to foreign markets. The effect of tillage becomes doubly exhaustive when by artificial means we stimulate vegetation without furnishing all the materials required for the growing plants. Such is the effect of many special manures, which while they supply certain elements, enable the plants to remove the others more rapidly from the soil. A partial exhaustion of the soil results likewise from repeated crops of the same kind; for the element of which the cereals require the largest quantity are taken in smaller proportions by green crops, and reciprocally, so that by judicious alterations the balance between the different mineral ingredients of the soil is preserved,

One of the great problems in scientific agriculture is to supply to the soil with the ammonia and the mineral matters necessary to support an abundant vegetation, and to obtain from various sources these different elements at prices which will permit of their being economically made use of. Nowhere but in the manure of the stable and farm-yard can we find combined all the fertilizing elements required, but several of them may be very cheaply procured. Thus lime and magnesia are abundant in the shape of marl and lime-stones; soda is readily obtained, together with chlorine, in common salt; while gypsum or plaster of Paris supplies at a low price both sulphuric acid and lime. Potash when wanting may be supplied to the soil by wood-ashes, but phosphoric acid and ammonia are less easily obtained and command higher prices.

An abundant supply of phosphate of lime is found in bones, which when dried contain from 50.0 to 60.0 p. c. of mineral matter, consisting of phos-