

RECORDS OF OTHER NORTHERN SPECIES IN OUR COLLECTION.

Melanophila drummondii Kirby.

Yukon Territory, lat. $62^{\circ} 31' 63^{\circ} 06'$, long. $137^{\circ} 30' 139^{\circ} 30'$, 1916, D. D. Cairnes, collector; near Sixtymile river, along 141st meridian, Yukon Territory, 1907, T. P. Reilly, collector, 2 specimens. Abundant throughout British Columbia, extending south into the United States and northward into the Yukon and Alaska. I have taken a series of thirteen near Lesser Slave lake in Alberta, and rarely in the east (Fort Conlonge, Que., in *Pinus strobus*).

Chrysobothris trinervia Kirby.

Near Sixtymile river, along 141st Meridian, Yukon Territory, 1907, Thos. P. Reilly, collector, 1 specimen.

There are two specimens in our collection from Fort Yukon, Alaska. Represented in our collection from Halifax, N.S., across Canada to Alaska and the Yukon.

Buprestis nuttalli Kirby.

Stewart river, Yukon Territory, 1909, D. H. Nelles, collector, 1 specimen.

There are only two small reddish-yellow spots on each elytron and the posterior of these almost obsolete on the right side; the spots on the outer faint but distinct on abdominal segments and coxae. Represented in our collections from Quebec, Ontario, Alberta, southern British Columbia, Alaska, and the Yukon.

Dicerca prolongata Lec.

Fort Chipewyan, Alta., 15-VI-1914, lake Athabaska, Alta., near shore on portage to lake One, 29-VI-14, F. Harper, collector, 1 specimen.

Dicerca tenebrica Ky.

Stewart river, Yukon Territory, D. H. Nelles, collector, 1909; Yukon Territory, latitude $62^{\circ} 31' 63^{\circ} 06' N.$, long. $137^{\circ} 30' 139^{\circ} 30' W.$, D. D. Cairnes, collector; lake Athabasca, Alta., near shore on portage to lake One, 29-VI-14, F. Harper, collector, 1 specimen; Dawson, Yukon Territory, 20-VI-08, 1 specimen.

Poecilnota, undes. sp.

Stewart river, Yukon Territory, D. H. Nelles collector, 1909.

Length, 17 mm.; black above, eyes red, venter black with coppery reflections, apparently distinct, 1 specimen.