

were covered with white membrane. He was injected September 19th. By Saturday, September 23rd, sores were all healing and were undergoing changes already described. By September 27th ulcers were all healed. There were a few tubercles left on the top of the shoulder. He was unable to take second injection. He returned January 4th, 1906, and reported that his cough had left him and that he had gained five pounds. The skin of the arm had broken down where the tubercles were left after first injection, also, a spot on the back of the neck. He was injected, and when seen January 10th the ulcers had healed again, and the skin had become soft and pliable. In a spot on the back of the neck the tubercles were seen to have undergone the usual change. He was injected January 26th, and at present, February 10th, 1906, skin lesion is healed.

Case VI.—Ernest L. came to the Montreal Dispensary, April 28th, 1905, complaining of a skin disease of several years duration, cough, purulent expectoration, and emaciation. There were three ulcers, one on arm, one on forearm, and one on back of right hand, near ulnar side. They were typical tubercular ulcers. For the last seven years there has been a certain amount of healing and then breaking down.

On examination of lungs dulness in right supraclavicular and infraclavicular regions was found, broncho-vesicular breathing, and sub-mucous rales. Sputum contained tubercle bacilli.

April 29th injected with iodoform gr. xxiv. There was a marked change in the ulcers. All three took on a healthy action. He was injected May 18th, and June 28th, 1905. After the last injection the sore on upper arm and forearm had healed completely. The one on the back of the hand was practically healed, only a few tubercles being left. The cough had almost ceased and patient's general health was very much better. He did not appear for treatment again.

Tuberculosis of the Lungs.

Case VII.—Mrs. K., aged 30, came to the Montreal Dispensary, March, 1905, complaining of cough, difficulty in deglutition, a hoarseness and rapid loss of flesh.

Family History.—Two sisters have died of consumption. Present illness began three years ago, December, 1902, with slight cough followed by a hæmorrhage from the lungs. During the winter patient had a hoarseness, cough and expectoration, but when the summer came the hoarseness cleared up. The cough continued up to October, 1904, when patient caught a severe cold. Cough got worse, and hoarseness returned. The patient at this time was pregnant, and was confined