

pens, R. Br.; *Cypripedium parviflorum*, Salisb.; *C. pubescens*, Willd.; *Triglochin palustre*, L.; *Eriophorum alpinum*, L., *Rhynchospora alba*, Vahl.; *Oryzopsis asperifolia*, Michx.; *Zizania aquatica*, L.; *Cystopteris bulbifera*, Bernh.; *Woodsia ilvensis*, R. Br.; *Aspideum cristatum*, Swartz; *Habenaria lacera*, R. Br. (common between Au Lac and Port Elgin); *H. viridis*, R. Br., var. *bracteata*, Reich., Petitcodiac and Havelock—Brittain; *Cuscuta Gronovii*, Willd., Nashwaaksis and St. Stephen, not rare about St. John, also at Lake Utopia. *Papaver dubium*, L., and *Calendula officinalis* have been produced from seeds self sown in gardens at St. Stephen for many years—Vroom; *Calamintha clinopodium*, Benth., *Nitella flexilis*, Agardh, are probably not rare.

G. U. HAY,
J. VROOM,
R. CHALMERS,

Committee on Botany.

ST. JOHN, January, 1884.

ADDITIONAL LIST OF PLANTS—1883.

- 30a. *GLAUCIUM LUTEUM*, Scop. Horn-Poppy. Ballast, St. John, Sep., G. U. Hay.
- 34a. *NASTURTIUM OFFICINALE*, R. Br. True Water-Cress. Hillsborough, Aug., J. Brittain.
42. *SISYMBRIUM OFFICINALE*, Scop. Spreading rapidly at St. Stephen, J. Vroom.
67. *SAPONARIA OFFICINALIS*, L. Along shore Long Reach, Saint John River, Sep., Hay.
- 77a. *Stellaria crassifolia*, Ehrh. Thick-leaved Chick-weed. Shediack, July, Brittain.
79. *S. humifusa*, Rottboell. Shediack, July, Brittain.
82. *Sagina procumbens*, L. The apetalous form (*S. apetala*, L.) occurs at St. Stephen, Vroom.

87b. C
88b. M
127. A
130a. O
166a. R
191a. P
209a. B
213. O
231a. V
232a. V. d
239a. C
241a. V
248a. A
249a. A. sa
270a. S
277a. X
277b. H
Isl