

Expanse 31 mil. Ohio (Mr. Drury).

*Botis socialis*, n. s.

Fore wings triangulate, widening outwardly more than usual. Ground color pale yellow washed with red, especially beyond the outer line, where a broad, diffuse, blackish subterminal band forms the most prominent marking of the wing and appears purplish from overlying bright scales. Transverse lines blackish, linear, trembled. Stigmata small, both solid. Outer line not strongly indented below median vein. Fringes pale, silky, very faintly interlined. The terminal edge of the wing is narrowly red. In one specimen the red shades are more or less absent, leaving the subterminal shade blackish. Hind wings pale yellow, with the subterminal broad shade more or less obviously continued; fringes pale. A fine mesial line. Beneath very pale sericeous yellow, with the subterminal shade and fragments of the exterior line repeated. On primaries discal marks repeated. On fore wings in one specimen the fringe is a little fuscous above. Body yellowish, paler beneath.

Expanse 25 mil. Canada, Mr. Saunders; Buffalo, Mr. Zesch.

*Botis allectalis*, n. s.

♂. Size of *communis*, but with more pointed primaries and longer abdomen. Gray over fuscous, with a pale ochreous discal patch on the median space surrounding the stigmata. These latter distinct, solid, dark fuscous, the oblique orbicular probably sometimes with paler centre, as it is faintly so on one wing. The lunate reniform followed by a fuscous shade margining the ochreous patch outwardly. T. p. line denticulate with a whitish included shade, setting out the line, not much indented below median vein. The gray scales overlay the fuscous, and when the wing is rubbed are first lost. The costa to t. p. line is ochrey fuscous. Hind wings translucent fuscous with pale line and darker borders. Head and appendages ochrey fuscous. Beneath paler with stigmata and exterior common line apparent.

Expanse 23 mil. Belfrage (No. 445), May 12, Bosque Co., Texas.