

inclined to think that this species may be confounded in some collections with *P. bilineata*, examples of which from Nova Scotia are in my collection.

---

### NOTE ON THECLA AUGUSTUS.

BY REV. THOMAS W. FYLES, SOUTH QUEBEC.

I beg to record the capture, by myself, in the neighborhood of Berger-ville, Province of Quebec, of two specimens of *Thecla Augustus*. One of them was taken on the 6th, and the other on the 8th of June. I am indebted to Mr. W. H. Edwards for the identification of the insects.

A figure of *Thecla Augustus* is given by Harris in his work on insects injurious to vegetation, page 279. As he gives no description of the insect, the following may not be unwelcome to some of the readers of the CANADIAN ENTOMOLOGIST:—

Expanse of wings 1 inch. Colour above, umber-brown, darker along the costa, and at the base in fore-wings. At the centre, in the fore-wings, there is a rust-red tinge or blush; and at the anal angle in the hind-wings there is an indistinct spot of the same color. The under-side of the fore-wings is of a lighter shade than the upper. Beyond the centre of the wing is a wavy transverse dark line. The hind-wings on the under-side have a basal patch of dark umber irregularly bordered. The antennæ are ringed black and white.

---

### STRAY NOTES ON MYRMELEONIDÆ, PART 2.

BY DR. H. A. HAGEN, CAMBRIDGE, MASS.

(Continued from page 136.)

#### 3. *Acanthaclisis Texana* Hagen.

I have a male and two females from Carrizo Spring, Dimmit Co., Western Texas, just near the frontier of Mexico.

Length of body, male, with app., 50 m.m.; female, 45. Length with wings, 65 m.m. Exp. al., 118 to 120.

Very similar in shape and color to *A. Americana*. After long consideration I believe them to be different species, until by a larger material