following contour of the margin, and above these, near end of discoidal space, are two slight dashes also. These markings are almost too indistinct to warrant mention.

Hind margins of both wings have a slight fringe of ground colour.

Under side of fore wing dark brown, but lighter and less rich than upper surface. The suggestions of dark markings are repeated. Inner marginal area a little lighter than balance of wing.

Under side of hind wing the same dark brown, the lighter markings of upper surface being repeated. The basal area has a few indistinct mottlings.

A coloured plate of a species like this is absolutely necessary for anything approaching accurate identification, and a plate will be duly published at a later date.

It may resemble Adelyodes castalus, Hew., but that has three subapical light spots, which do not appear at all in this, and on close inspection of the scale distribution on under side of hind wing, considerable difference will be noticed.

Taken in November, 1899.

Plestia Kikkawai, sp. nov.—Habitat : Suapure, Venezuela. Expanse, 1.25 inches.

Head, thorax and abdomen above dark brown, with a few grayish hairs. Circles of grayish hairs at base of each segment of abdomen. There is a white dot at base of each antenna. Antennæ dark brown, tip beneath gray. Legs nearly white.

Upper side of fore wing dark brown, with nine white spots or markings, giving the wing almost a mottled appearance. The largest of these is at centre of discoidal space, strongly concaved on marginal side. Below this, in the next two lower interspaces, are two white spots. This row of spots is edged with very dark brown on basal side. Near the end of discoidal space, near costa, is a small white spot. There are two larger white spots or markings in the first and second submedian interspaces, both of them more or less concaved on marginal side. There are three apical white dots. Below them, near lower angle, is a well-defined marking of very dark.

Upper side of hind wing is dark brown. Costa and portion of discoidal space close to base crossed by a heavy white band. Below this, near centre of wing, is a prominent white spot, with a smaller one adjoining it. A band of very light brown extends on each side of these spots towards upper angle and inner margin respectively, giving the