

WHICH APPEAR TO HAVE BEEN PERIRENAL IN ORIGIN.

TERMINATION.	RELATIONSHIP TO KIDNEY.	NATURE OF TUMOUR.	WEIGHT.	REMARKS.
No operation.	In connection with capsule of right kidney.	Some parts firmer than others; appeared like fat.	—	"Ut steatomatis refferret materiam."
—	Left kidney behind and to inner side of tumour.	Pure Lipoma; very vascular.	315 grms.	Tumour only found at autopsy, 2 to 3 times size of kidney; soft and fluctuating; capsule of kidney continued on to tumour
Laparotomy; peritonitis; d. in 48 hours.	Right kidney involved in mass.	Lipoma.	—	Even after removal of main mass 12 to 20 lbs. of the growth were removed at autopsy.
Laparotomy; fistula from descending colon; recovery.	Left kidney adherent to border of tumour in a kind of hilum.	Fibro-lipoma oedematousum.	37½ lbs, (German)	Puncture attempted several times; states origin of growth from capsule: gives good diagram, but calls growth a "fibroma."
Death from respiratory troubles.	R. kidney buried 1½ inches in growth.	Myxo-lipoma.	41½ lbs,	Describes case as a myxoma which had undergone fatty degeneration.
Laparotomy; d. 2 days later.	"Started from left kidney."	Lipoma, not fully described.	Enormous.	Puncture negative; diagnosis echinococcus cysts; spleen also adherent.
Laparotomy: d. shortly after.	Left kidney completely imbedded in upper part of tumour.	Myxo-lipoma.	Size of man's head.	Echinococcus diagnosed.
Laparotomy; col-lapse.	Right kidney carried four fingers breadth forward by growth.	Fibro-lipoma myxomatodes.	Size of man's head.	Solid ovarian tumour diagnosed.
Laparotomy; col-lapse.	Left kidney imbedded in the growth.	Myxo-lipoma.	27,400 grms.	Puncture negative; gives good diagram; points out peri-renal origin.
—	Assoc'd with capsule of left kidney.	Fibro-lipoma.	11 kilos.	Diagnosed as multilocular cyst of ovary.
Laparotomy; recovery: recurrence.	Left kidney removed with tumour.	In part lipoma, in part myxoma fibromatosum.	10 kilo (with kidney).	Diagnosed ovarian cyst; recurrence after 1 year; inoperable.
Laparotomy; recovery.	Degenerated kidney, side not stated, thought to be within tumour.	Lipoma.	6,600 grms,	No signs of kidney, its vessels, or ureter seen during the operation; in middle of tumour a mass of fatty degenerated tissue shape of kidney (?)
Pyæmia and asphyxia; death.	Left kidney imbedded in hinder surface and in part atrophied.	Myxo-lipoma becoming sarcomatous in one area.	14 lbs.	Puncture negative; diagnosis retroperitoneal tumour confirmed by exploratory laparotomy.
Light cold; d. from exhaustion.	Left kidney imbedded in hinder surface and atrophied.	Myxo-lipoma.	41½ lbs.	—