

NEW PRINCIPLE FOR THE FAT ASSIMILATION OF HYDROLEINE "HYDRATED OIL."

"HYDROLEINE" may be described as partially digested oil, which will nourish and produce increase in weight, in those cases where oils or fats, not so treated, are difficult or impossible to digest. In CONSUMPTION and other WASTING DISEASES, the most prominent symptom is *emaciation*, of which the first is the starvation of the fatty tissues of the body, including the brain and nerves. This tendency to emaciation and loss of weight is arrested by the regular use of HYDROLEINE. The ordinary so-called emulsions of Cod Liver Oil and other fats, *whether pancreatised or not*, merely remain in the form of a coarse mechanical mixture for a short time after agitation. The digestion of oil, having in no sense been artificially produced, still devolves upon those functional powers, the deficiency of which is the most prominent symptoms in these cases.

A great misconception as to the real characteristics of a true pancreatic emulsion has been entertained by many, and but few appear to have studied the different aspects presented by such an emulsion as is produced on fat by the energetic action of pure soluble pancreatin, as contrasted with the coarse mechanical mixtures of oil or fat and water, which are commonly supposed to represent this function of fermentative digestion.

Some seem to think that if a bottle of oil is shaken up with the compounds sold as the active principle of the pancreas, and a yellowish cloud is diffused for a time through the oil, an emulsion has been obtained. So it has, but not the true pancreatic emulsion, which forms an integral portion of the process by which fats are digested and assimilated. From the unvarying result of many hundred trials with the pure, active principles of healthy pancreatic fluid, taken at the time of digestion, I am perfectly convinced that no valuable result has been attained, unless the emulsion formed is as highly refractive of light as milk. The color may vary, according to the oil or fat used, from a far whiter fluid than the densest milk to the opacity and color of Devonshire cream, but unless at least the equivalent of the density of the best milk is produced in oil, when a third of water is held in suspension, no real pancreatic emulsion has been formed.

The mere mechanical mixture formed by common pancreatin is rarely better or more persistent than may be produced by rubbing up oil or fat with a solution of mucilage, or by a warm application of dissolved gelatin, shaken with oil until it becomes cold.

The first essential towards the digestion of fats or oils in the human body is that it shall assume the state of the very finest and most permanent emulsion, and this is only known to be attained when the oil and water is perfectly opaque, from the minuteness of the globules. This is the first function of the pancreatic emulsifying principle, and by this alone can we be certain that it possesses its proper fermentative activity.—Prof. Bartlett's Treatise.

(HYDRATED OIL)

HYDROLEINE

(WATER AND OIL.)

The efficacy of this Preparation is NOT CONFINED to cases of CONSUMPTION, as from its valuable tonic effect on the nervous system, in addition to its special stimulating action on the organs concerned in the production of Fat in the body, it causes a *natural increase in weight in persons of naturally thin habit, who do not present any evidence of disease.*

The principles upon which this discovery is based have been described in a treatise on "THE DIGESTION AND ASSIMILATION OF FATS IN THE HUMAN BODY," by H. C. BARTLETT, PH. D., F.C.S., and the experiments which were made, together with cases illustrating the effect of Hydrated Oil in practice, are concisely stated in a treatise on "CONSUMPTION AND WASTING DISEASES," by G. OVEREND DREWRY, M.D., of London.

In these treatises, the Chemistry and Physiology of the Digestion of Fats and Oils is made clear, not only by the description of a large number of experiments scientifically conducted, but by cases in which the deductions are most fully borne out by the results.

Copies of these valuable works will be sent free on application.

FORMULA OF HYDROLEINE.

Each dose of two teaspoonsful, equal to 120 drops, contains:

Pure Oil.....	80 m (drops.)
Distilled Water.....	35 "
Soluble Pancreatin.....	5 grains.
Soda.....	$\frac{1}{2}$ "
Boric Acid.....	$\frac{1}{2}$ "
Hyocholeic Acid.....	1-20 "

DOSE.—Two teaspoonsful alone, or mixed with twice the quality of soft water, to be taken thrice daily with meals.

Unlike the ordinary preparation of Cod-Liver Oil, it produces no unpleasant eructation or sense of nausea, and should be taken in such very much smaller doses, according to the directions, as will insure its complete assimilation; this, at the same time, renders its use economical in the highest degree.

Upon application from any of the Medical Faculty, I will be pleased to forward samples, which will substantiate the claims made for Maltpepsyn, and I hope for your assistance in this my endeavour to introduce a good preparation at a low price.