

POLYPETALOUS EXOGENS.

RANUNCULACEÆ—Crow-foot Family.

CLEMATIS, L. Traveller's Joy. Virgin's Bower.

1. C. VERTICILLARIS, D.C. (Mauve-flowered Clematis).

Atragene Americana. Sims.

Creeping over rocks and low shrubs. King's and other mountains, Chelsea, P.Q. Not uncommon. May—4.

2. C. VIRGINIANA. (Common Virgin's-Bower.)

Borders of streams. Aug.—1.

ANEMONE, L. Wind Flower.

12. A. CYLINDRICA, Gray (Long-fruited Anemone).

Dry pastures. Ju.—2.

13. VIRGINIANA, L.

Rocky, open woods. There are two forms of this plant: (i) with greenish flowers like the preceding, but always with much shorter heads of fruit; and (ii) with white flowers almost as large as those of No. 15. From this latter, however, it is easily distinguished by the woolly seeds. (i) Common; (ii) on the rocks round Lake Flora, Hull, P.Q. Ju.—2. (B.)

15. A. DICHOTOMA, L. (Round-headed Anemone).

A. Pennsylvanica, L.

Along river margins and in low meadows. Ju.—2. (B.)

17. A. HEPATICA, L. (Liverwort. Blue May-flower).

Hepatica triloba, Chaix.

Dry, rocky woods. One of our earliest flowers; not quite as common as No. 18. Ap.—3 (B.)

18. A. ACUTILOBA, Lawson ("May-flower").

Dry, shady woods. Ap.—3 (B.)

THALICTRUM, Tourn. Meadow-rue.

20. T. DIOICUM, L. (Early Meadow-rue).

River banks and low woods. Ju.—1. (B.)