

52. *HALOPORPHYRUS VIOLA*, Goode & Bean. Fishing banks off the coast (U. S. F. C.)
53. *ONOS (RHINONEMUS) CIMBRIUS (L.)* Goode & Bean. *Motella cimbria*, Gunth. Fishing banks off the coast (U. S. F. C.)
54. *BROSMIUS BROSMÆ (Mull.)* White. Cusk. Common.
55. *AMMODYTES AMERICANUS*, De Kay. Sand Eel. Common; burying in the sand at ebb of tide, and going in schools at high water.

Fam. PLEURONECTIDÆ.

56. *HIPPOGLOSSUS VULGARIS*, Flem. Halibut. Very common on the fishing banks off the coast.
57. *HIPPOGLOSSOIDES PLATESSOIDES (Fabr.) Gill.* Arctic Flounder. *Hippoglossoides limandoides*, Gunth. Not uncommon. Of two specimens forwarded by the Rev. J. Ambrose from St. Margaret's Bay, the largest measured twenty-two inches in length.
58. *PSEUDOPLEURONECTES AMERICANUS (Walb.) Gill.* Flounder. *Platessa plana*, Stor. Very common in shore waters. Described by the author, Trans. N. S. Inst. Nat. Sc., Vol. I, Pt. I., p. 51.
59. *LIMANDA FERRUGINEA (Storer.)* Goode & Bean. *Pleuronectes ferrugineus*, Gunth.
60. *PLATYSOMATICHTHYS HIPPOGLOSSOIDES (Walb.)* Goode & Bean. Turbot. *Hippoglossus groenlandicus*, Gunth. Occasional specimens are brought in from the northern fishing banks, but it is more common off Newfoundland. It is a very oily fish when cooked.
61. *GLYPTOCEPHALUS CYNOGLOSSUS (L.) Gill.* *Pleuronectes cynoglossus*, Gunth. La Have fishing bank (U.S.F.C.)

Fam. SILURIDÆ.

62. *AMIURUS CATUS (L.) Gill.* Not common. This fish is very tenacious of life, for a specimen survived being carried wrapped up in paper in a pocket for two hours.

Fam. SALMONIDÆ.

63. *SALMO SALAR, L.* Salmon. More abundant some years