

H. tessellaris Sm. Abb.

- 1797—Sm. Abb.*, Ins. Ga., II., 149, pl. 75, *Phalæna*.
 1816—Hbn., Verzeichniss, 170, *Halisidota*.
 1837—Geyer, Zutr., No. 470, ff. 939-940, *Halisidota*.
 1833—Harris, Cat. Ins., Mass., 592, *Arctia*.
 1841—Harris*, Rept. Ins., Mass., 260, *Lophocampa*.
 1856—Wlk., C. B. Mus., Lep. Het., III., 732, *Halisidota*.
 1860—Clem., Proc. Ac. Nat. Sci., Phil., XII, 534, *Halisidota*.
 1862—Morris, Synopsis App., 348, *Halisidota*.
 1862—Harris*, Inj. Insects, 364, *Lophocampa*.
 1863—Saund.*, Syn. Can. Arct., 19, *Halisidota*.
 1864—Walsh*, Proc. Bost. Soc. N. H., IX., 289, *Halisidota*.
 1870—Walsh*, Am. Ent., I., 205, *Lophocampa*.
 1873—Stretch*, Zyg. & Bomb., 87, 137, pl. VI., f. 9, ♀, *Halisidota*.
 1882—Grote, New List, 16, *Halisidota tessellata*.

antiphola Walsh.

- 1864—Walsh*, Proc. Bost. Soc. N. H., IX., 288, *Halisidota*.
 1864—Walsh*, Proc. Ent. Soc., Phil., III., 412, 413, *Halisidota*.
 1864—Grt., Proc. Ent. Soc., Phil., III., 536, *pr. syn.*
 1865—Walsh, Proc. Ent. Soc., Phil., V., 197, *pr. syn.*

var. harrisii Walsh.

- 1864—Walsh, Proc. Ent. Soc., Phil., III., 430, *Halisidota*.
 1865—Walsh, Proc. Ent. Soc., Phil., V., 197, *pr. var.*
 1870—Walsh, Am. Ent., I., 205, *Lophocampa*,
 1873—Stretch, Zyg. and Bomb., 137, *pr. var.*

Habitat—Canada to Texas; Illinois, Indiana, Missouri.

Food-plants—Oak, sycamore, beech, hornbeam, plane.

H. trigona Grt.

- 1879—Grt., No. Am. Ent., I., 46, *Halisidota*.
 1881—Grt., Trans. Kans. Ac. Sci., VII., 64, *Halisidota*.
 Habitat—New Mexico.

Genus EUHALESIDOTA, Grt.

- 1865, Grt., Proc. Ent. Soc., Phil., V., 243.

Head moderate; eyes large, not prominent; tongue of medium length, corneous; palpi short, scarcely exceeding the front; terminal joint very short; antennæ long, in the male heavily pectinated; legs increasing in length posteriorly; spurs normal in number and moderate in length.