

it does not differ as far as I can see. The species may be known by the discoloured purplish fuscous head and thorax, the dark color extending on the shoulders of the fore wings. It has been collected in Texas by Belfrage in March and July. The primaries are more pointed in *orgiae* and this species is more faintly colored when compared with its congeners.

The species may be thus catalogued :

*Erotyla* Hübn.

*sulphuralis* Linn. Europe.

*Spragueia* Grote.

*onagrus* Guen., 2, 205. Florida.

*Type* *leo* Guen., 2, 205. Alabama.

*plumbifimbriata* Grote, C. E., ix., 68. Texas.

*dama* Guen., 2, 205. Alabama, Texas.

*guttlata* Grote, C. E., vii., 225. Texas.

*tortricina* Zeller, Beitr., i., 15. Texas.

*apicella* Grote, Trans. Am. Ent. Soc., iv., 21. Alabama, Texas.

*truncatula* Zeller, Beitr., i., 3.

*Fruva* Grote.

*Type* *fasciatella* Grote, C. E., vii., 225. Texas.

*obsoleta* Grote, C. E., ix., 69. Illinois, Texas.

*Xanthoptera* Guen.

*Type* *nigrofimbria* Guen., 2, 241. N. Y., Ala., Texas.

*semiflava* Guen., 2, 241. Texas.

*Exyra* Grote.

*Type* *semicrocea* Guen., 2, 241. Georgia.

*Ridingsii* Riley, Trans. St. Louis Acad., iii., 240. Georgia.

*nigrocaput* Morrison, Proc. Bost. Soc. N. H., 17, 153.

*fax* Grote, Trans. Am. Ent. Soc., 295. Georgia.

*Rolandiana* Grote, Psyche, ii., 38. Mass.