

quantity so as to form an indistinct, pale, irregular fascia at the beginning of the ciliae; at the apex the ochreous brown again prevails over the white. At the base of the costa the ochreous fuscous is very narrow. Posterior wings purplish fuscous; under surface and legs whitish, the legs marked with brownish on their anterior surfaces. *Al. ex.* $\frac{1}{2}$ inch.

L. rufocristatella. N. sp.

White; antennae annulate with brown. There is a small reddish ochreous tuft on the disc before the middle of the primaries, behind which the wing is suffused with reddish ochreous; a larger tuft nearly opposite to the first one, and nearer to the dorsal margin, of the same hue with the first one, and another just within the dorsal margin opposite the beginning of the ciliae. Between this last tuft and the costa, and thence to the apex, the wing is suffused with reddish ochreous, sparingly dusted with fuscous. *Al. ex.* $\frac{1}{2}$ inch. Season, July.

L. ignotilisella. N. sp.

No raised tufts on the wings. Silvery white, the costal margin about the middle dusted with pale purple and ochreous scales: an irregular streak or sinus of spots, which are connected by ochreous brown dusting, begins before the middle of the disc, and extends backwards, spreading over the apical part of the wing. *Al. ex.* $\frac{1}{2}$ inch. Season, September.

L. albocapitella. N. sp.

Head, thorax and base of the dorsal margin of the wings pure snowy white; antennae purplish fuscous, iridescent; palpi white, stained externally with pale purplish fuscous; the white patch at the base of the dorsal margin of the primaries is posteriorly narrowly margined with dark brown, containing a small raised tuft on the margin. The remainder of the primaries are of an indescribable hue, composed of grayish brown ochreous and white scales intermixed, the former hue predominating, or rather, it is brown streaked with ochreous and sprinkled with white; there are three small tufts of dark brown raised scales, each anteriorly margined with white, the white followed by a margin of dark brown; two of them are on the disc before the middle, the other further back and nearer the dorsal margin, and behind it is a fourth tuft also near the dorsal margin; there is a short dark brown streak in the apical part of the wing, within the dorsal margin, and a small dark brown spot about the end of the disc. *Al. ex.* $\frac{1}{2}$ inch. Season, September.