

my material, and it is not clear that Mr. Morrison has identified a species more nearly resembling Hübner's figure than the ordinary identification of *tritona*.

*Apatela grisea* (Walk).

I have examined Mr. Walker's types in the British Museum, and I made the note: Fore wings like *tritona*; hind wings white. Professor Bélanger sent me a species which I considered to be this of Walker's. It now appears to me that Mr. Morrison has altered my determination and described the moth as a new species. I think that, until Mr. Walker's species is more satisfactorily identified, my own determination should not be interfered with.

I have since identified several of the species marked unknown to me in the "List" of 1874. A shorter compilation of the "List" is now published. It is no proof that the species is unknown now, that it was at that time. For instance, I have since identified *Schinia gracilentia*, and it seems to me that it is the same as *oleagina* Morr.; in my specimen the subterminal space is a little darker than terminal, as somewhat exaggeratedly shown in Hübner's figure; there is a faint discal discoloration beneath and traces of a subterminal line; the hind wings above are hardly "rosy" along exterior border. My specimen is undoubtedly the same species as Mr. Morrison's, and also came from my old school-friend Mr. Graef. I come to the conclusion that *oleagina* is not a var., but a synonym of *gracilentia*.

*Apatela dentata*, n. s.

♂. This is allied to *tritona* and *grisea*, but is a smaller species wanting all the black dashes. The ground color is blackish, shaded over with whitish. The lines black, single, denticulate. The claviform is indicated by a slight black mark. Orbicular obsolete. Reniform whitish, rounded, with its outward edge black-lined and shaded. T. a. line running in a little on median vein. T. p. line inaugurated above the reniform, running well outwardly, denticulate throughout its length. Subterminal line hardly apparent; a blackish shade over median nervules on the gray terminal space. Fringes gray preceded by blackish interspaceal markings. Hind wings fuscous, lighter towards the base, with indistinct line. Beneath much paler, irrorate, without discal marks and with a common shaded line.

Expanse 32 m. m. Quebec, Mr. Bowles.