

Vaginal examination revealed a slightly thickened os and slight endocervicitis with erosions of the cervix. Cod liver oil, malt extract, hypophosphites, and aromatics, in combination, 25 per cent. of each, were given freely during the intervals between the menstrual periods and for three days before the expected menstruation Ergoapiol (Smith) was given in capsules, one being given three times daily until the discharge ceased. At the fourth period after the beginning of the treatment she was relieved of all her symptoms, and was free from pain and fetor during menstruation. Locally, tincture of iodine and occasionally tampons of ichthyol and glycerine were applied. The cure was permanent and in every way satisfactory.

MUSCULAR SORENESS AND RHEUMATISM DUE TO GRIP.

In speaking of the treatment of articular rheumatism, Hobart A. Hare, M.D., Professor of Therapeutics in the Jefferson Medical College and Editor of *The Therapeutic Gazette*, says: "Any substance possessing strong antipyretic power must be of value under such circumstances." He further notes that the analgesic power of the coal-tar products "must exert a powerful influence for good." The lowering of the fever, no doubt, quiets the system and removes the delirium which accompanies the hyperpyrexia, while freedom from pain saves an immense amount of wear, and places the patient in a better condition for recovery. The researches of Guttman show conclusively that these products possess a direct anti-rheumatic influence, and among those remedies antikamnia stands pre-eminent as an analgesic and antipyretic. Hare, in the last edition of his *Practical Therapeutics* says: "Salol renders the intestinal canal antiseptic." This is much needed in the treatment of rheumatism. In short, the value of salol in rheumatic conditions is so well understood and appreciated that further comment is unnecessary. The statements of Professors Hare and Guttman are so well known and to the point and have been verified so often, that we are not surprised that the wide-awake manufacturers placed "Antikamnia and Salol Tablets" on the market. Each one of these tablets contains two and one-half grains of antikamnia and two and one-half grains of salol. The proper proportion of the ingredients is evidenced by the popularity of the tablets in all rheumatic conditions and particularly in that condition of muscular soreness which accompanies and follows the grip. The Antikamnia Chemical Company, St. Louis, Mo., will send samples to physicians on application. Please mention this journal.
