

quite excited, delirious and headstrong, determined to get out of bed. After sweating for an hour, temp. dropped to 102, pulse, 96.

Dec. 11th. Fifteen days after my first visit morning temp. normal. One stool slightly tinted with bile, odor normal; evening temp., 102, pulse, 100, no dulness.

Dec. 13th. Bowels moved by enema. Three stools well stained with bile, temp. 102, pulse 100. Zinc sulpho-carb. stopped; ordered Watson's chlorine mixture, 2 drams, with 2 gr quinine every four hours; bowels to be moved by enema second day.

After this the temperature and pulse gradually fell, reaching normal on the 27th of Dec., 32 days after the onset. Patient objected to chlorine water and the following tablet was substituted: strychnine, 1-60 gr., ipecac, 1-10 gr., capsicum, 1-4 gr., ext. rhubarb, 1-4 gr., ext. gentian, 1-2 gr., soda 2 gr., three times a day before meals; light diet.

Patient got out of bed Jan. 1st, and gradually regained health and strength.

Jan. 24th, 1906. After driving over the rough roads a distance of about ten miles and eating a hearty dinner, she again complained of pain over the stomach and right side, with vomiting. When I visited her at 8 p.m., temperature 101, pulse 90, and bowels not moved for two days. I gave her a dose of calomel to be followed by mag. sulph. in the morning, and left salicylate of soda mixture to be taken four times a day after the bowels moved. Her temperature was normal the next day, and she has remained well ever since.

CASE 2.—July 27th, 1905, while passing was called to see J. R., age 8. His mother told me he had been drowsy and not feeling well for two or three days. During the previous 24 hours bowels had been moved three times, the last two passages being quite fetid; tongue covered with a whitish fur, not red on tips or edges; temp. 100, pulse 80; slight tenderness or gurgling in the right iliac fossa, no headache, pupils normal.

Gave sodium salicylate, 3 gr., with ext. glycyrrhiza every three hours; milk diet, and told them to report his condition if not well in a few days.

Saw nothing more of him and heard nothing from him until three weeks later he came to my office, August 17th, when he had a well-marked jaundice, and on examination I found the lower border of the liver on a level with the umbilicus, the ribs being somewhat raised upwards, the whole abdomen bulged forward and very prominent. On palpation there was no doubt of there being an enlarged liver, not markedly tender; his temp. was normal, pulse 60. On enquiry I learned that after my previous visit, though ordered milk diet, he had been eating anything and everything that he wished.